

MATHEMATICS – BASIC

SOLUTIONS

SAMPLE QUESTION PAPER

CBSE Class X Examination

6

Section ‘A’

[1 × 20 = 20]

1. (d) 2. (b) 3. (a) 4. (d)
 5. (d) 6. (b) 7. (d) 8. (b)
 9. (c) 10. (c)

Fill in the blanks

11. $k = 8$

Explanation

Given equation is $x^2 - 3x + k - 10 = 0$

Here $a = 1, b = -3$ and $c = k - 10$

$$\text{Product of roots} = \frac{c}{a}$$

$$-2 = \frac{k-10}{1}$$

$$\Rightarrow k = 8$$

12. $k = -11$

Explanation

Putting $x = 3$ in $3x^2(k-1)x + 9 = 0$

$$3(3)^2 + (k-1)(3) + 9 = 0$$

$$27 + 3k - 3 + 9 = 0$$

$$3k = -33$$

$$k = -11$$

13. $x = 5, 3$

Explanation

Given equation is $x^2 - 8x + 15 = 0$

$$x^2 - 5x - 3x + 15 = 0$$

$$x(x-5) - 3(x-5) = 0$$

$$(x-5)(x-3) = 0$$

$$x = 5, 3$$

14. $c = 6$

Explanation

$$x^2 - 5x + c = 0$$

Here $a = 1, b = -5$ and $c = c$

$$\therefore \alpha + \beta = \frac{-b}{a} = 5$$

$$\alpha + \beta = 5$$

$$\text{Adding } \frac{\alpha - \beta = 1}{2\alpha = 6} \text{ or } \alpha = 3, \beta = 2$$

$$\text{Now product of roots } \alpha\beta = \frac{c}{a}$$

$$3 \times 2 = \frac{c}{1}$$

$$c = 6$$

15. $k = 6$

Explanation


For infinitely many solutions

$$\frac{2}{4} = \frac{3}{k} = \frac{5}{10}$$

$$\Rightarrow k = 6$$

Answer the following.

16. Applying pythagoras theorem.



$$PO^2 = 3^2 + 4^2$$


$$= 25$$

$$\therefore PQ = 5 \text{ units}$$

$$\begin{aligned} 17. \cos(40 + \theta) - \sin(50 - \theta) \\ &= \cos[90 - (50 - \theta)] - \sin(50 - \theta) \\ &= \sin(50 - \theta) - \sin(50 - \theta) \\ &= 0 \end{aligned}$$

18. $\sin 60^\circ \cdot \cos 30^\circ + \cos 60^\circ \cdot \sin 30^\circ$

$$\begin{aligned} &= \frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2} + \frac{1}{2} \times \frac{1}{2} \\ &= \frac{3}{4} + \frac{1}{4} \\ &= 1 \end{aligned}$$



Given $\angle AOB = 130$

Also $\angle APB + \angle AOB = 180^\circ$

$\therefore \angle APB = 180^\circ - 130^\circ = 50^\circ$

20. $\sec \theta + \tan \theta = x$... (i)

Rationalising

$$\begin{aligned} &\frac{(\sec \theta + \tan \theta)}{1} \times \frac{(\sec \theta - \tan \theta)}{(\sec \theta - \tan \theta)} = x \\ &\Rightarrow \frac{\sec^2 \theta - \tan^2 \theta}{\sec \theta - \tan \theta} = \frac{x}{1} \\ &\qquad\qquad\qquad (\sec^2 \theta - \tan^2 x = 1) \\ &\Rightarrow \sec \theta - \tan \theta = \frac{1}{x} \qquad\qquad\qquad \dots \text{(ii)} \end{aligned}$$

from (i) and (ii)

$\sec \theta + \tan \theta = x$

$\sec \theta - \tan \theta = \frac{1}{x}$

Adding $2 \sec \theta = x + \frac{1}{x}$

$$2 \sec \theta = \frac{x^2 + 1}{x}$$

$$\sec \theta = \frac{x^2 + 1}{2x}$$

Section ‘B’

21. $\text{LCM}(p, q) = a^3b^3$

[½]

$$\text{HCF}(p, q) = a^2b$$

[½]

$$\text{LCM}(p, q) \times \text{HCF}(p, q) = a^5b^4$$

$$= (a^2b^3)(a^3b) = pq \quad [1]$$

22. (i) Let the cost of one book be ₹ x and that of one pen be ₹ y . Then,

$$\text{Cost of 5 books} + \text{cost of 7 pen} = ₹ 79$$

$$\Rightarrow 5x + 7y = 79 \quad \dots(i)$$

$$\text{Cost of 7 books} + \text{Cost of 5 pens} = ₹ 77$$

$$\Rightarrow 7x + 5y = 77 \quad \dots(ii)$$

Adding (i) and (ii),

$$5x + 7y = 79$$

$$+ 7x + 5y = 77$$

$$12x + 12y = 156$$

$$12(x + y) = 156$$

$$x + y = \frac{157}{12} = 13 \quad [½]$$

$$100(x + y) = 13 \times 100 = ₹ 1300$$

∴ 100 books and 100 pen will cost ₹ 1300. [1]

(ii) Pair of linear equation in two variables. [½]

23. First 8 multiples of 3

$$3, 6, 9, 12, 15, 18, 21, 24$$

$$S = 3 + 6 + 9 + 12 + 15 + 18 + 21 + 24$$

These numbers are in A.P.

where $a = 3, d = 3$ and $n = 8$


$$S_n = \frac{n}{2}[2a + (n-1)d]$$

$$S_8 = \frac{8}{2}[2 \times 3 + (8-1)3]$$

$$S_8 = 4[6 + 21]$$

$$S_8 = 4 \times 27 = 108$$

∴ Thus, sum of first 8 multiples of 3 of 108. [1]



P divides AB 1 : 2


∴ Coordinates of P are : (-1, 0) [1]

∴ Q is mid-point of PB

∴ Coordinates of Q are : (-4, 2) [1]

[CBSE Marking Scheme, 2016]

CBSE, Topper's Solution, 2016



25. Total number of all possible outcomes = 20

Numbers multiple of 3 are : 3, 6, 9, 12, 15, 18

Numbers multiple of 7 are : 7, 14 [1]

Numbers multiple of 3 or 7 are : 3, 6, 7, 9, 12, 14, 15, 18

Let A be the event that the number on the drawn ticket is a multiple of 3 or 7.

Then, the number of favourable outcomes = 8

$\therefore P$ (that the drawn ticket is a multiple of

$$3 \text{ or } 7) = P(A) = \frac{8}{20} = \frac{2}{5} \quad [1]$$

26. Possibilities are HH, HT, TH, TT [1]

$$P(\text{HH or TT}) = \frac{2}{4} = \frac{1}{2} \quad [1]$$

Section 'C'

27. Let us consider two numbers 225 and 60

$$\begin{aligned} \text{We obtain} \quad 225 &= 60 \times 3 + 45 & [1] \\ &\because 225 > 62 \end{aligned}$$

$$60 = 45 \times 1 + 15$$

[$\because 60 > 45$]

$$45 = 15 \times 3 + 0$$

[$\because 45 > 15$]

Thus, HCF (225, 60) = 15 [1]

Also, $225 \times 5 - 10x = 15$

$$\Rightarrow 10x = 1125 - 15$$

$$\Rightarrow x = \frac{1110}{10} = 111 \quad [1]$$

28. Let $\alpha = 1$ and $\beta = -3$.

Sum of zeros = $(\alpha + \beta) = 1 + (-3) = -2$.

Product of zeros = $\alpha\beta = 1 \times (-3) = -3$. [1/2]


So, the required polynomial is

$$\begin{aligned} x^2 - (\alpha + \beta) + \alpha\beta &= x^2 - (-2)x + (-3) \\ &= x^2 + 2x - 3. \quad [1] \end{aligned}$$

$$\text{Sum of zeros} = -2 = \frac{-2}{1} = \frac{-(\text{coefficient of } x)}{(\text{coefficient of } x^2)},$$

$$\text{Product of zeros} = -3 \quad [1]$$

$$= \frac{-3}{1} = \frac{\text{constant term}}{\text{coefficient of } x^2}. \quad [1\frac{1}{2}]$$



ABCD is a parallelogram

\therefore diagonals AC and BD bisect each other
Therefore

Mid point of BD is same as mid point of AC

$$\Rightarrow \left(\frac{a+1}{2}, \frac{2}{2} \right) = \left(\frac{-2+4}{2}, \frac{b+1}{2} \right)$$


$$\Rightarrow \frac{a+1}{2} = 1 \quad \text{and} \quad \frac{b+1}{2} = 1$$

$\Rightarrow a = 1, b = 1$. Therefore
length of sides are $\sqrt{10}$ units each.

[1+1+1] [CBSE Marking Scheme, 2016]

Expert's Solution:

We know that diagonals of parallelogram
bisect each other.



\therefore Midpoint of diagonal AC

$$\left(\frac{-2+4}{2}, \frac{1+b}{2} \right) = \left(1, \frac{1+b}{2} \right) \quad [1\frac{1}{2}]$$

Mid-point of diagonal BD

$$\left(\frac{a+1}{2}, \frac{0+2}{2} \right) = \left(\frac{a+1}{2}, 1 \right) \quad [1\frac{1}{2}]$$

Mid point of diagonal AC \cong mid point of
diagonal BD

$$1 = \frac{a+1}{2} \quad \text{and} \quad \frac{1+b}{2} = 1$$

$$2 = a + 1 \quad \text{and} \quad 1 + b = 2 \\ \therefore \quad a = 1 \quad \text{and} \quad b = 1 \quad [1]$$

$$AB = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$= \sqrt{(1+2)^2 + (0-1)^2}$$

$$AB = \sqrt{9+1} = \sqrt{10} \text{ unit}$$

$$BC = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$= \sqrt{(4-1)^2 + (1-0)^2}$$

$$BC = \sqrt{9+1} = \sqrt{10} \text{ unit}$$

ABCD is a parallelogram (Given)

$$AB = CD = \sqrt{10} \text{ unit} \quad [1\frac{1}{2}]$$

$$BC = AD = \sqrt{10} \text{ unit} \quad [1\frac{1}{2}]$$

OR

$$\text{Given,} \quad AP = \frac{3}{7} AB$$


$$\Rightarrow AP = \frac{3}{7} (AB + BP)$$

$$\Rightarrow 7AP = 3AP + 3BP \quad [1]$$

$$\Rightarrow 4AP = 3BP$$

$$\Rightarrow \frac{AP}{BP} = \frac{3}{4}$$

$$\Rightarrow AP : BP = 3 : 4 \quad [1]$$



Now, coordinates of P

$$= \left(\frac{3 \times 2 + 4 \times (-2)}{3+4}, \frac{3 \times (-4) + 4 \times (-2)}{3+4} \right)$$

[by internal section formula]


$$= \left(\frac{6-8}{7}, \frac{-12-8}{7} \right) = \left(\frac{-2}{7}, \frac{-20}{7} \right) \quad [1]$$

30. Let AB be the vertical stick and let AC be its shadow.

Then, AB = 0.15 m and AC = 0.12 m.

Let DE be the vertical tower and let DF be its shadow.

Then, DF = 50 m. Let DE = x metres.



Now, in $\triangle BAC$ and $\triangle EDF$, we have:

$$\angle BAC = \angle EDF = 90^\circ$$

$$\angle ACB = \angle DFE \quad [1]$$


[angular elevation of the sun at the same time]

$\therefore \triangle BAC \sim \triangle EDF$.

$$\begin{aligned} \Rightarrow \frac{AB}{DE} &= \frac{AC}{DF} \Rightarrow \frac{0.15}{x} = \frac{0.12}{50} \\ \Rightarrow x &= \frac{(0.15 \times 50)}{0.12} = 62.5. \quad [1] \end{aligned}$$

Hence, the height of the tower is 62.5 m.

OR



Distance covered by first aeroplane due North after two hours = $500 \times 2 = 1,000$ km. $\quad [\frac{1}{2}]$

Distance covered by second aeroplane due East after two hours = $650 \times 2 = 1,300$ km. $\quad [\frac{1}{2}]$

Distance between two aeroplane after 2 hours

$$\begin{aligned} NE &= \sqrt{ON^2 + OE^2} \\ &= \sqrt{(1000)^2 + (1300)^2} \\ &= \sqrt{1000000 + 1690000} \\ &= \sqrt{2690000} \\ &= 1640.12 \text{ km.} \end{aligned}$$

31. LHS = $(\operatorname{cosec} \theta - \cot \theta)^2$

$$= \left(\frac{1}{\sin \theta} - \frac{\cos \theta}{\sin \theta} \right)^2 = \left(\frac{1 - \cos \theta}{\sin \theta} \right)^2 \quad [1]$$

$$= \frac{(1 - \cos \theta)^2}{\sin^2 \theta} = \frac{(1 - \cos \theta)(1 - \cos \theta)}{(1 - \cos^2 \theta)} \quad [1]$$

$$[\because \sin^2 \theta = 1 - \cos^2 \theta]$$

$$= \frac{(1 - \cos \theta)}{(1 + \cos \theta)} = \text{RHS.} \quad [1]$$

$\therefore \text{LHS} = \text{RHS.}$

OR

$$\text{LHS} = \frac{\cos A}{1 - \tan A} + \frac{\sin A}{1 - \cot A}$$

$$= \frac{\cos A}{1 - \left(\frac{\sin A}{\cos A} \right)} + \frac{\sin A}{1 - \left(\frac{\cos A}{\sin A} \right)}$$

$$= \frac{\cos^2 A}{\cos A - \sin A} + \frac{\sin^2 A}{\sin A - \cos A}$$

$$= \frac{\cos^2 A}{\cos A - \sin A} - \frac{\sin^2 A}{\cos A - \sin A}$$


$$= \frac{\cos^2 A - \sin^2 A}{\cos A - \sin A}$$

$$= \frac{(\cos A - \sin A)(\cos A + \sin A)}{(\cos A - \sin A)}$$

$$= \cos A + \sin A$$

$$= \sin A + \cos A = \text{RHS}$$

32. We know, $AC = r$
 In $\triangle ACB$, $BC^2 = AC^2 + AB^2$
 $\Rightarrow BC = AC\sqrt{2}$
 $\quad (\because AB = AC)$
 $\Rightarrow BC = r\sqrt{2} \quad [1]$



Required area = $ar(\Delta ACB) + ar(\text{semicircle on BC as diameter}) - ar(\text{quadrant ABPC})$

$$= \frac{1}{2} \times r \times r + \frac{1}{2} \times \pi \times \left(\frac{r\sqrt{2}}{2} \right)^2 - \frac{1}{4} \pi r^2 \quad [1]$$

$$= \frac{r^2}{2} + \frac{\pi r^2}{4} - \frac{\pi r^2}{4}$$

$$= \frac{r^2}{2} = \frac{196}{2} \text{ cm}^2 = 98 \text{ cm}^2 \quad [1]$$

[CBSE Marking Scheme, 2018]

$$\therefore \text{Value of sphere} = \frac{4}{3} \times \frac{22}{7} \times r^3 \quad [1]$$

$$= 504 \times \frac{1}{3} \times \frac{22}{7} \times \frac{35}{20} \times \frac{35}{20} \times 3$$

$$= \frac{4}{3} \times \frac{22}{7} \times r^3 \quad [1]$$

$r = 10.5 \text{ cm. } \therefore \text{diameter} = 21 \text{ cm}$

$$\text{Surface area} = 4 \times \frac{22}{7} \times \frac{21}{2} \times \frac{21}{2}$$

$$= 1386 \text{ cm}^2 \quad [1]$$

[CBSE Marking Scheme, 2015]

33. Volume of metal in 504 cones

$$= 504 \times \frac{1}{3} \times \frac{22}{7} \times \frac{35}{20} \times \frac{35}{20} \times 3 \text{ cm}$$

CBSE Topper's Solution, 2015

(17) Diameter of a cone = 3.5 cm
 $\Rightarrow \text{radius}(R) = \frac{3.5}{2} \text{ cm}$

Height of cone = H = 3 cm

Here, 504 cones are melted to form a sphere.

Let the radius of sphere = R

Now, Vol^m of sphere = Vol^m of 504 cones

or, Vol^m of $\frac{4}{3} \pi R^3 = 504 \times \frac{1}{3} \times \frac{22}{7} \times 3 \times \frac{3.5}{2} \times \frac{3.5}{2} \times 3$

or, $R^3 = \frac{126 \times 3 \times 3.5 \times 3.5}{4 \times 160} = \frac{7 \times 7 \times 7 \times 9}{16 \times 8}$

or, $R = \sqrt[3]{\frac{7 \times 7 \times 7 \times 3 \times 3 \times 3}{2 \times 2 \times 2}} = \frac{7 \times 3}{2} = \frac{21}{2} \text{ cm}$

$\therefore \text{Diameter} = 2R = 2 \times \frac{21}{2} = 21 \text{ cm}$

S.A of hemispherical canal = $4 \pi R^2$
 $= 4 \pi \times \frac{21}{2} \times \frac{21}{2} = \frac{22}{7} \times 21 \times 21$
 $= 66 \times 21$
 $= 1386 \text{ cm}^2$

OR

Let the area that can be irrigated in 30 minutes be A m².

Water flowing in canal in 30 minutes

$$= \left(10,000 \times \frac{1}{2} \right) \text{ m} = 5000 \text{ m}$$

Volume of water flowing out in 30 minutes
 $= (5000 \times 6 \times 1.5) \text{ m}^3 = 45000 \text{ m}^3 \quad \dots(i) \quad [1]$

Volume of water required to irrigate the field

$$= A \times \frac{8}{100} \text{ m}^3 \quad \dots(ii)$$

Equating (i) and (ii), we get

$$\begin{aligned} A \times \frac{8}{100} &= 45000 \\ A &= 562500 \text{ m}^2. \end{aligned}$$

[CBSE Marking Scheme, 2018]

34.

| Salary (in thousand ₹) | No. of persons (f) | cf |
|------------------------|------------------------|------|
| 5–10 | 49 | 49 |
| 10–15 | 133 | 182 |
| 15–20 | 63 | 245 |
| 20–25 | 15 | 260 |
| 25–30 | 6 | 266 |
| 30–35 | 7 | 273 |
| 35–40 | 4 | 277 |
| 40–45 | 2 | 279 |
| 45–50 | 1 | 280 |

[2]

$$\frac{N}{2} = \frac{280}{2} = 140$$

Median class is 10–15

$$\begin{aligned} \text{Median} &= l + \frac{h}{f} \left(\frac{N}{2} - C \right) \\ &= 10 + \frac{5}{133} (140 - 49) = 10 + \frac{5 \times 91}{133} = 13.42 \end{aligned}$$

Median salary is ₹ 13.42 thousand or ₹ 13420
(approx) [1]

[CBSE Marking Scheme, 2016]

Section 'D'

$$35. x + 2y = 5 \text{ or, } y = \frac{5-x}{2}$$

| | | | |
|---|---|---|---|
| x | 1 | 3 | 5 |
| y | 2 | 1 | 0 |

[1]

$$2x - 3y = -4$$


$$\text{Or, } y = \frac{2x+4}{3} \quad [1]$$

| | | | |
|---|---|---|----|
| x | 1 | 4 | -2 |
| y | 2 | 4 | 0 |

[1]

lines meet x -axis at (5, 0) and (-2, 0)
respectively.

Graph [2]

**OR**

Let the time taken by the tap to fill the tank of smaller diameter = x hours

\therefore Time taken by the tap of larger diameter to fill the tank = $(x - 9)$ hours

\therefore Work done by the tap of smaller diameter

$$\text{in one hour} = \frac{1}{x} \quad [1/2]$$

and the work done by the tap of larger

$$\text{diameter in one hour} = \frac{1}{x-9} \quad [1/2]$$

Thus, the work done by the two taps together in 1 hour

$$\begin{aligned} &= \frac{1}{x} + \frac{1}{x-9} \\ &= \frac{x-9+x}{x(x-9)} = \frac{2x-9}{x(x-9)} \quad [1] \end{aligned}$$

The two tap together can fill the tank in $\frac{x(x-9)}{2x-9}$ hours.

According to the information,

$$\frac{x(x-9)}{2x-9} = 6 \text{ hours (given)}$$


$$\begin{aligned} &\Rightarrow x^2 - 9x = 12x - 54 \\ &\Rightarrow x^2 - 21x + 54 = 0 \quad [1] \\ &\Rightarrow x^2 - 3x - 18x + 54 = 0 \\ &\Rightarrow x(x-3) - 18(x-3) = 0 \\ &\Rightarrow (x-3)(x-18) = 0 \\ &\Rightarrow \text{Either } x-3 = 0 \\ &\text{or } x-18 = 0 \\ &\Rightarrow \text{Either } x = 3 \text{ or } x = 18 \quad [1] \end{aligned}$$

When $x = 3$, then the tap of smaller diameter can fill the tank in 3 hours and the tap of the larger diameter can fill the tank in $3 - 9 = -6$, which is rejected as time to fill the tank cannot be negative.

When $x = 18$, then the tap of smaller diameter can fill the tank in 18 hours and the tap of the larger diameter can fill the tank in $18 - 9 = 9$ hours.

- 36. Given:** A trapezium ABCD in which $AB \parallel DC$ and $AB = 2DC$. Its diagonals intersect each other at the point O.

To find: $\frac{\text{ar}(\Delta AOB)}{\text{ar}(\Delta COD)}$



Method of Solution: In ΔAOB and ΔCOD , we have:

$$\angle AOB = \angle COD \text{ (vert. opp. } \angle\text{s)}$$

$$\angle OAB = \angle OCD \text{ (corres. } \angle\text{s)}$$

$\therefore \Delta AOB \sim \Delta COD$ (by AA-similarity). [1]

We know that the ratio of the areas of two similar triangles is equal to the ratio of the squares of the corresponding sides.

$$\begin{aligned} \therefore \frac{\text{ar}(\Delta AOB)}{\text{ar}(\Delta COD)} &= \frac{AB^2}{DC^2} \\ &= \frac{(2 \times DC)^2}{DC^2} [\because AB = 2 \times DC] [1] \\ &= \frac{4 \times DC^2}{DC^2} = \frac{4}{1}. \end{aligned}$$

Hence, $\text{ar}(\Delta AOB) : \text{ar}(\Delta COD) = 4 : 1$. [1]

OR

Given : A ΔABC , in which AD is the median.

To prove : $AB^2 + AC^2 = 2(AD^2 + BD^2)$

Construction : Draw AE $\perp BC$.

Proof : In right ΔAEB , we have

$$AB^2 = AE^2 + BE^2 \quad \dots(i)$$

[By Pythagoras Theorem]

In right ΔAEC , we have

$$AC^2 = AE^2 + CE^2 \quad \dots(ii)$$

[By Pythagoras Theorem]


Adding (i) and (ii), we get

$$\begin{aligned} AB^2 + AC^2 &= (AE^2 + BE^2) \\ &\quad + (AE^2 + CE^2) \end{aligned}$$

$$\Rightarrow AB^2 + AC^2 = 2AE^2 + (BE^2 + CE^2) \quad \dots(iii) [1]$$

CBSE Topper's Solution, 2016

In right ΔAED , we have



[1]

$$AD^2 = AE^2 + ED^2$$

[By Pythagoras Theorem]

$$\Rightarrow AD^2 = AD^2 - ED^2 \quad \dots(iv)$$

From (iii) and (iv), we get

$$\begin{aligned} AB^2 + AC^2 &= 2(AD^2 - ED^2) + (BE^2 + CE^2) \\ &= 2AD^2 - 2ED^2 + BE^2 + CE^2 \\ &= 2AD^2 + (BE^2 - ED^2) + (CE^2 - ED^2) \\ &= 2AD^2 + (BE + ED). \end{aligned}$$

$$(BE - ED) + (CE + ED) (CE - ED)$$

$$= 2AD^2 + (BE + ED).BD$$

$$+ DE.(CE - ED)$$

$$= 2AD^2 + (BE + ED).BD$$

$$+ BD.(CE - ED) [1]$$

[$\because BD = CD$]

$$= 2AD^2 + BD.(BE + ED + CE - ED)$$

$$= 2AD^2 + BD.(BE + CE)$$

$$= 2AD^2 + BD.BC$$


$$= 2AD^2 + BD(2BD)$$

[$\because AD$ is median]


$$= 2(AD^2 + BD^2) \quad [1]$$

37. Correct Construction. [4]

[CBSE Marking Scheme, 2016]



38.



[1]

Let AB be the tower and ships are at points C and D.

$$\tan 45^\circ = \frac{AB}{BC}$$

$$\Rightarrow \frac{AB}{BC} = 1$$

$$\Rightarrow AB = BC \quad [1]$$

$$\text{Also, } \tan 30^\circ = \frac{1}{\sqrt{3}} = \frac{AB}{BC+CD}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{AB}{AB+CD} \quad [1]$$

$$\Rightarrow AB + CD = \sqrt{3}AB$$

$$\Rightarrow CD = AB(\sqrt{3}-1)$$

$$= 100 \times (1.732 - 1)$$

$$= 73.2 \text{ m} \quad [1]$$

[CBSE Marking Scheme, 2018]

39. Radius of the sphere = 3 cm

$$\text{Volume of the sphere} = \left(\frac{4}{3} \pi r^3 \right) \text{cm}^3$$

$$= \left(\frac{4}{3} \pi \times 3 \times 3 \times 3 \right) \text{cm}^3 = (36\pi) \text{ cm}^3. \quad [1]$$

Length of wire 36 cm.

Let the radius of the wire be r cm.

$$\text{Volume of the wire} = (\pi r^2 h) \text{ cm}^3$$

$$= (\pi r^2 \times 36) \text{ cm}^3 \quad [1]$$

But, volume of wire = volume of sphere

$$\Rightarrow 36\pi r^2 = 36\pi \quad [1]$$

$$\Rightarrow r^2 = 1 \quad [1]$$

$\Rightarrow r = 1$ [$\because r$ cannot be negative].

Hence, the radius of the wire is 1 cm.

40.

| C.I. | f | c.f. |
|-------|---|----------|
| 0–10 | 5 | 5 |
| 10–20 | x | $5+x$ |
| 20–30 | 6 | $11+x$ |
| 30–40 | y | $11+x+y$ |
| 40–50 | 6 | $17+x+y$ |
| 50–60 | 5 | $22+x+y$ |

[2]

Here from table, $N = 22 + x + y = 40$

$$\Rightarrow x + y = 18 \quad \dots(i)$$

Since, Median = 31, which lies between 30–40

$$\therefore \text{Median} = 30-40$$

$$\text{Median} = l + \left(\frac{\frac{N}{2} - c.f.}{f} \right) \times h$$

$$\Rightarrow 31 = 30 + \left[\frac{20 - (11+x)}{y} \right] \times 10$$

$$\Rightarrow 1 = \frac{(9-x) \times 10}{y}$$

$$\Rightarrow y = 90 - 10x$$

$$10x + y = 90 \quad \dots(ii) \quad [1]$$

On solving eqn. (i) and (ii), we get

$$x = 8 \text{ and } y = 10 \quad [1]$$

OR

$$\text{Modal class} = 60-80 \quad [1]$$

$$\therefore Z = l + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times h \quad [1]$$

Here, $l = 60$, $f_1 = 29$, $f_0 = 21$, $f_2 = 17$ and $h = 20$

$$\text{Mode} = 60 + \frac{29-21}{2 \times 29 - 21 - 17} \times 20 \quad [1]$$

$$= 60 + \frac{8}{58-38} \times 20$$

$$= 60 + 8 = 68 \quad [1]$$

MATHEMATICS – BASIC

SOLUTIONS

SAMPLE QUESTION PAPER

CBSE Class X Examination

7

Section 'A'

[1 × 20 = 20]

1. (b) 2. (c) 3. (c) 4. (d)
 5. (c) 6. (b) 7. (c) 8. (c)
 9. (b) 10. (b)

Fill in the blanks

11. $k = -\frac{1}{3}$


Explanation

If three points are collinear, then area of triangle formed by them is zero.

$$A = \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)]$$

$$O = \frac{1}{2} [k(3k - 1) + 3k(1 - 2k) + 3(2k - 3k)] = 0$$

$$O = 3k^2 - k + 3k - 6k^2 - 3k$$



$$\begin{aligned} -3k^2 - k &= 0 \\ -k[3k + 1] &= 0 \\ \Rightarrow k &= -\frac{1}{3} \end{aligned}$$

12. $AB = 6\sqrt{2}$ cm

Explanation

$AC = BC$ (Isosceles Δ) Applying Pythagoras theorem

$$AB^2 = AC^2 + BC^2$$

$$AB^2 = 6^2 + 6^2$$

$$AB^2 = 36 + 36$$

$$AB^2 = 2 \times 36$$

$$AB = 6\sqrt{2}$$
 cm

13. $\theta = 126^\circ$

Explanation

$$\text{Area of sector} = \frac{7}{20} \text{ Area of circle}$$

$$\frac{\theta}{360} \times \pi r^2 = \frac{7}{20} \pi r^2$$

$$\Rightarrow \theta = 126^\circ$$

14. $\frac{16}{9}$

Explanation

$$\text{Given: } \frac{V_1}{V_2} = \frac{64}{27}$$

$$\frac{\frac{4}{3}\pi r_1^3}{\frac{4}{3}\pi r_2^3} = \left(\frac{4}{3}\right)^3$$

$$\left(\frac{r_1}{r_2}\right)^3 = \left(\frac{4}{3}\right)^3$$

$$\therefore \frac{r_1}{r_2} = \frac{4}{3}$$

$$\frac{s_1}{s_2} = \frac{4\pi r_1^2}{4\pi r_2^2} = \left(\frac{r_1}{r_2}\right)^2 = \left(\frac{4}{3}\right)^2 = \frac{16}{9}$$

15. 0

Explanation

$$\operatorname{cosec}(75 + \theta) - \sec(15 - \theta)$$

$$= \operatorname{cosec}(90 - 15 + \theta) - \sec(15 - \theta)$$

$$= \operatorname{cosec}[90 - (15 - \theta)] - \sec(15 - \theta)$$

$$= \sec(15 - \theta) - \sec(15 - \theta) \\ = 0$$

Answer the following.

16.
$$\frac{1 + \tan^2 \theta}{1 + \cot^2 \theta} \quad \left[\text{As } 1 + \tan^2 \theta = \sec^2 \theta \right]$$

$$= \frac{\sec^2 \theta}{\operatorname{cosec}^2 \theta}$$

$$= \frac{1}{\cos^2 \theta} \times \frac{\sin^2 \theta}{1}$$

$$= \tan^2 \theta$$

17. $S_p = 2p^2 + 3p$

$$S_1 = 2 + 3 = 5 = a_1$$

$$S_2 = 8 + 6 = 14 = a_1 + a_2$$

$$\therefore a_2 = 9$$

$$\Rightarrow d = 9 - 5 = 4$$

18. Let $\alpha = -2$ and $\beta = 3$

Sum $\alpha + \beta = -2 + 3 = 1$

Product $\alpha\beta = (-2)(3) = -6$

\therefore Quadratic polynomial $= x^2 - 8x + P = 0$

$\Rightarrow x^2 - x - 6 = 0$

Now comparing $a + 1 = -1$

$\Rightarrow a = -2$

$b = -6$

19. The given equation $7x^2 - 12x + 18 = 0$

Here $a = 7, b = -12$ and $c = 18$

Sum of roots $= \frac{-b}{a} = \frac{12}{7}$

Product of roots $= \frac{c}{a} = \frac{18}{7}$

\therefore Ratio $= \frac{12}{18} = \frac{2}{3}$

20. $x^2 - 4x + 3 = 0$

$x^2 - 3x - x + 3 = 0$

$x(x - 3) - 1(x - 3) = 0$

$$(x - 3)(x - 1) = 0 \\ x = 3, 1$$

Section 'B'

21. $\frac{1717}{2^2 \times 5^3} = \frac{1717 \times 2}{2^3 \times 5^3} = \frac{3434}{(10)^3} = \frac{3434}{1000} = 3.434$ [2]

22. Given, system of equations is

$$2x + 3y = 7$$

$$\text{and } (a - 1)x + (a + 1)y = 3a + 1$$

On comparing these equations with $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$,

$$\text{we get, } a_1 = 2, b_1 = 3, c_1 = -7$$

$$\text{and } a_2 = (a - 1), b_2 = (a + 1), c_2 = -(3a + 1)$$

For parallel lines (no solution), we have

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2} = \frac{2}{a-1} = \frac{3}{a+1} \neq \frac{-7}{-(3a+1)}$$

$$\Rightarrow 2(a + 1) = 3(a - 1)$$

$$\Rightarrow 2a + 2 = 3a - 3$$

$$\Rightarrow a = 5$$

$$\text{Also, } \frac{3}{a+1} \neq \frac{7}{3a+1}$$

$$\Rightarrow 9a + 3 \neq 7a + 7 \quad \therefore a = 5$$

$$2a \neq 4$$

$$a \neq 2$$

23. Give A.P. is $-6, -9, -12, -15 \dots$

Here, $a = -6$,

Common difference (d) $= -9 + 6 = -3$ [1]

\therefore 11th term of the given A.P. $-6, -9, -12, 15 \dots$

$$T_n = a + (n - 1)d$$

$$T_{11} = -6 + (11 - 1)(-3)$$

$$I = -6 - 30$$

$$= -36$$

24. $PA^2 = PB^2$

$$\Rightarrow (x - 5)^2 + (y - 1)^2$$

$$= (x + 1)^2 + (y - 5)^2$$

$$\Rightarrow 12x = 8y$$

$$\Rightarrow 3x = 2y$$

CBSE Topper's Answer, 2017

$$\begin{aligned}
 PA &= PB \\
 \therefore PA^2 &= PB^2 \\
 \text{by distance formula,} \\
 (5-x)^2 + (1-y)^2 &= (-1-x)^2 + (5-y)^2 \\
 \Rightarrow (5-x)^2 + (1-y)^2 &= (1+x)^2 + (5-y)^2 \\
 25 - 10x + x^2 + 1 - 2y + y^2 &= 1 + 2x + x^2 + 25 - 10y + y^2 \\
 -10x - 2y &= 2x - 10y \\
 \frac{8y}{4(2y)} &= \frac{12x}{4(3x)} \\
 \therefore 3x &= 2y \\
 \text{Hence, proved.}
 \end{aligned}$$

25. Here, total number of playing cards = 52
 Number of red cards and queens = $26 + 2 = 28$ [½]
 ∴ Number of cards other than red cards and queens = $52 - 28 = 24$ [½]
 Required probability of getting neither a red card nor a queen = $\frac{24}{52} = \frac{6}{13}$ [1]

26. Total no. of pens = 144
 Defective one = 20

$$\begin{aligned}
 \text{Good ones} &= 144 - 20 = 124 \\
 \text{(i) Probability of purchasing pen} &= \frac{124}{144} = \frac{31}{36} \\
 \text{(ii) Probability of not purchasing pen} &= \frac{24}{144} = \frac{5}{36} \quad [1]
 \end{aligned}$$

Section 'C'

27. $1 \times 2 \times 3 \times 4 \times 5 \times 6 \times 7 + 5 = 5(1 \times 2 \times 3 \times 4 \times 5 \times 6 \times 7 + 1)$ [1]
 $= 5(1008 + 1)$
 $= 5(1009)$
 $=$ a composite number
 $[\because \text{product of two factors}]$ [1]

28. Let α, β, γ be the zeroes of the given polynomial $f(x)$, such that

$$\begin{aligned}
 \alpha + \beta &= 0 \\
 \text{sum of zeroes} &= \frac{-\text{coefficient of } x^2}{\text{coefficient of } x^3} = -4 \\
 \Rightarrow \alpha + \beta + \gamma &= \frac{-4}{1} \\
 \Rightarrow 0 + \gamma &= -4 \quad [\alpha + \beta = 0] \\
 \gamma &= -4 \quad [1]
 \end{aligned}$$

Product of zeroes

$$\begin{aligned}
 &= \frac{-\text{constant term}}{\text{coefficient of } x^3} \Rightarrow \alpha\beta\gamma = \frac{-(-36)}{1} = 36 \\
 \therefore \alpha(-\alpha)(-4) &= 36 \quad [1] \\
 \therefore 4\alpha^2 &= 36 \Rightarrow \alpha^2 = 9 \Rightarrow \alpha = \pm 3 \\
 \text{Now, } \alpha + \beta &= 0, \beta = \pm 3 \\
 \text{Hence zeroes are } 3, -3, -4. \quad [1]
 \end{aligned}$$

29. Here

$$\begin{aligned}
 \frac{1}{2} \{(k+1)(-3+k) + 4(-k-1) + 7(4)\} &= 6 \quad [1] \\
 \Rightarrow k^2 - 6k + 9 &= 0 \quad [1] \\
 \text{Solving to get } k &= 3 \quad [1]
 \end{aligned}$$

[CBSE Marking Scheme, 2015]

CBSE Topper's Answer, 2015

Given, vertices of a triangle be
 $A(k+1, 1)$, $B(4, -3)$ and $C(7, -k)$

Given, $\text{ar}(\Delta ABC) = 6 \text{ sq. units}$

Then, $x_1 = k+1$, $y_1 = 1$
 $x_2 = 4$, $y_2 = -3$
 $x_3 = 7$, $y_3 = -k$

Now, $\text{ar}(\Delta ABC) = \frac{1}{2} |x_1(y_2-y_3) + x_2(y_3-y_1) + x_3(y_1-y_2)|$

Or, $6 = \frac{1}{2} |(k+1)(-3+k) + 4(-k-1) + 7(1+3)|$

Or, $12 = (k+1)(k-3) - 4k - 4 + 28$

Or, $12 = k^2 - 2k - 3 - 4k + 24$

Or, $k^2 - 6k + 21 - 12 = 0$

Or, $k^2 - 6k + 9 = 0$, which is a quad. eqn.

Or, $k^2 - 3k - 3k + 9 = 0$

Or, $k(k-3) - 3(k-3) = 0$

Or, $(k-3)(k-3) = 0$

Or, $k-3 = 0$ Or, $k = 3$

∴ $k = 3$

OR

Diagonals of a parallelogram bisect each other.

So, mid-point of $AC = \text{mid-point of } BD$

$$\text{i.e., } \left(\frac{1+k}{2}, \frac{2-2}{2} \right) = \left(\frac{2-4}{2}, \frac{3-3}{2} \right)$$

$$\Rightarrow \left(\frac{1+k}{2}, 0 \right) = (-1, 0)$$

$$\text{i.e., } \frac{(1+k)}{2} = -1$$

$$\text{i.e., } k = -3$$

Now $\text{ar}(ABCD) = 2 \text{ Area of } \Delta ABD$

$$= 2 \times \frac{1}{2} \times [1(6) + 2(-1) - 4(-5)]$$

$$= 24 \text{ sq. units.} \quad [1]$$

$$AB = \sqrt{(2-1)^2 + (3+2)^2} = \sqrt{26} \text{ units}$$

$$\text{ar}(ABCD) = \text{base} \times \text{height} = AB \times h$$

$$\text{So, } 24 = \sqrt{26} \times h$$

$$\text{So, } h = \frac{24}{\sqrt{26}} \text{ units} \quad [1]$$


30. Given: ΔABC and ΔDBC are drawn on the same hypotenuse BC and on the same side of BC . Also, AC and BD intersect at P .

To Prove: $AP \times PC = BP \times PD$ **Proof:** In ΔBAP and ΔCDP , we have:

$$\angle BAP = \angle CPD = 90^\circ$$

$$\angle BPA = \angle CPD \text{ (ver. opp. } \angle\text{s)}$$

$$\therefore \Delta BAP \sim \Delta CDP \text{ [by AA-similarity]}$$




$$\begin{aligned}\therefore \frac{AP}{DP} &= \frac{BP}{CP} \\ \Rightarrow AP \times CP &= BP \times DP \\ \Rightarrow AP \times PC &= BP \times PD \\ \text{Hence, } AP \times PC &= BP \times PD.\end{aligned}$$

OR**Proof:**

In quadrilateral ABCD,

$$\begin{aligned}\frac{AO}{BO} &= \frac{CO}{DO} && \text{(Given)} \\ \text{or, } \frac{AO}{CO} &= \frac{BO}{DO} && \dots(i) [1]\end{aligned}$$

Draw EO \parallel AB on

[1]

In $\triangle ABD$, $EO \parallel AB$ (By construction)

$$\therefore \frac{AE}{ED} = \frac{BO}{DO} \quad \text{(By BPT)...(ii)}$$

From eqns. (i) and (ii),

$$\frac{AE}{ED} = \frac{AO}{CO}$$

$$\text{In } \triangle ADC, \frac{AE}{ED} = \frac{AO}{CO}$$

or, $EO \parallel DC$ (Converse of BPT) $EO \parallel AB$ (Construction) $\therefore AB \parallel DC$ or, In quad. ABCD, $AB \parallel DC$

or, ABCD is a trapezium.

[1]


Hence Proved

31. LHS = $\operatorname{cosec}^2 \theta - \tan^2 (90^\circ - \theta)$

$$\begin{aligned}&= \frac{1}{\sin^2 \theta} - \frac{\sin^2 (90^\circ - \theta)}{\cos^2 (90^\circ - \theta)} \\ &= \frac{1}{\sin^2 \theta} - \frac{\sin^2 (90^\circ - \theta)}{\sin^2 \theta} \quad [1]\end{aligned}$$

$$\begin{aligned}&= \frac{1}{\sin^2 \theta} - \frac{\cos^2 \theta}{\sin^2 \theta} \\ &= \frac{1 - \cos^2 \theta}{\sin^2 \theta} \quad [1] \\ &= \frac{\sin^2 \theta}{\sin^2 \theta} \\ &= 1 \\ &= \sin^2 \theta + \cos^2 \theta \\ &= \sin^2 \theta + \sin^2 (90^\circ - \theta) \quad [1] \\ &= \text{RHS}\end{aligned}$$

32. Let O be the centre of a circle of radius 5.6 cm, and let OACBO be its sector with perimeter 27.2 cm.

Then, $PA + PB + \text{arc } AB = 27.2 \text{ cm}$ [1]

$$\Rightarrow 5.6 \text{ cm} + 5.6 \text{ cm} + \text{arc } ACB = 27.2 \text{ cm}$$

$$\Rightarrow \text{arc } ACB = 16 \text{ cm.} \quad [1]$$


Area of the sector OACBO

$$\begin{aligned}&= \left(\frac{1}{2} \times \text{radius} \times \text{arc} \right) \text{sq units} \\ &= \left(\frac{1}{2} \times 5.6 \times 16 \right) \text{cm}^2 = 44.8 \text{ cm}^2. \quad [1]\end{aligned}$$

33. Heap of rice is cone shaped:


Diameter of its base = 24 m

Height of cone = 3.5 m



$$\begin{aligned}\text{In } \triangle OAP, OA^2 &= OP^2 + AP^2 \\ &= (12)^2 + (3.5)^2\end{aligned}$$

$$\begin{aligned}
 &= 144 + 12.25 = 156.25 \\
 \therefore OA(l) &= \sqrt{156.25} = 12.5 \text{ cm} \quad [1] \\
 \text{Volume of cone} &= \frac{1}{3}\pi r^2 h \\
 &= \frac{1}{3} \times \frac{22}{7} \times 12 \times 12 \times 3.5 \\
 &= 22 \times 4 \times 12 \times 0.5 \\
 &= 22 \times 2 \times 12 = 528 \text{ m}^3 \quad [1] \\
 \therefore \text{Volume of cone} &= 528 \text{ m}^3 \\
 \text{Canvas cloth is required to just cover the heap} \\
 &= \text{C.S.A. of cone} = \pi r l \\
 &= \frac{22}{7} \times 12 \times 12.5 = \frac{22 \times 150.0}{7} \\
 &= \frac{3300}{7} \\
 \text{Canvas cloth} &= 471.43 \text{ m}^2 \text{ (Approx)} \quad [1]
 \end{aligned}$$

OR

CBSE Topper's Answer, 2016

$$\begin{aligned}
 &\text{Radius of cylinder as well as conical part} = \frac{3}{2} \text{ m.} \\
 &\text{Height of cylinder, } h = 2.1 \text{ m} \\
 &\text{Slant height of cone, } l = 2.8 \text{ m.} \\
 &\text{Total canvas required} = 2\pi rh + \pi rl \\
 &\Rightarrow \frac{22}{7} \times \frac{3}{2} [4.2 + 2.8] \text{ m}^2 \\
 &\Rightarrow \frac{22}{7} \times \frac{3}{2} \times 7.0 \text{ m}^2 = 33 \text{ m}^2 \\
 &\text{Total cost @ ₹ } 500/\text{m}^2 = \boxed{\text{₹ } 16,500}
 \end{aligned}$$

| Height | Frequency | c.f. |
|---------|--------------|------------|
| 100–140 | 22 | 22 |
| 140–180 | 14 | 36 |
| 180–220 | 18 | 54 |
| 220–260 | 16 | 70 |
| 260–300 | 30 | 100 |
| | Total | 100 |

[1]

$$\begin{aligned}
 \text{Here, } N &= 100 \\
 \Rightarrow \text{Median} &= \frac{N}{2} \text{th term} \\
 &= \frac{100}{2} = 50 \text{th term} \\
 \text{So, Median Class} &= 180 - 220
 \end{aligned}$$

$$\text{Median} = l + \left(\frac{\frac{N}{2} - c.f.}{f} \right) \times h \quad [1]$$

$$= 180 + \left(\frac{50-36}{18} \right) \times 40 = 180 + \frac{14 \times 40}{18}$$

$$= 180 + 31.11$$

$$\text{Median} = 211.11 \quad [1]$$

Section 'D'

35. $(x-1)^2 + (2x+1)^2 = 2(2x+1)(x-1) \quad [1]$
 $\Rightarrow x^2 + 1 - 2x + 4x^2 + 1 + 4x = 4x^2 - 4x + 2x - 2 \quad [1]$
 $\Rightarrow x^2 + 4x + 4 = 0 \quad [1]$
 $\Rightarrow (x+2)^2 = 0 \quad [1]$
 $\Rightarrow x = -2 \quad [1]$

[CBSE Marking Scheme, 2017]

CBSE Topper's Answer, 2017

Let $\frac{x-1}{2x+1}$ be y ,

$$y + \frac{1}{y} = 2$$

$$y^2 + 1 = 2y$$

$$y^2 - 2y + 1 = 0$$

$$y^2 - y - y + 1 = 0$$

$$y(y-1) - 1(y-1) = 0$$

$$(y-1)^2 = 0$$

$$\therefore y = 1 \text{ or } -1.$$

Now, $\frac{x-1}{2x+1} = 1 \quad \text{or} \quad \frac{x-1}{2x+1} = -1$

$$x-1 = 2x+1$$

$$-2 = x$$

$$\therefore x = -2 \quad \text{or} \quad 1$$

$\therefore x = -2$

OR

Let the original speed be x km/h.
 Then, reduced speed = $(x - 400)$ km/h
 The duration of flight at original speed

$$= \left(\frac{6000}{x} \right) \text{hours} \quad \left[\because \text{time} = \frac{\text{distance}}{\text{speed}} \right]$$

The duration of flight at reduced speed

$$= \left(\frac{6000}{x-400} \right) \text{hours} \quad \dots(1)$$

Since, the difference between these two duration is $\frac{1}{2}$ hours.

$$\therefore \frac{6000}{x-400} - \frac{6000}{x} = \frac{1}{2} \quad [1]$$

$$\Rightarrow \frac{6000x - 6000(x-400)}{x(x-400)} = \frac{1}{12000}$$

$$\Rightarrow \frac{6000\{x-(x-400)\}}{x(x-400)} = \frac{1}{12000}$$

$$\Rightarrow \frac{x-x+400}{x(x-400)} = \frac{1}{12000}$$

$$\Rightarrow \frac{400}{x(x-400)} = \frac{1}{12000}$$

$$\Rightarrow x(x-400) = 4800000$$

$$\Rightarrow x^2 - 400x - 4800000 = 0 \quad [1]$$

$$\Rightarrow x^2 - 2400x + 2000x - 4800000 = 0$$

[by factorisation]

$$\Rightarrow x(x-2400) + 2000(x-2400) = 0$$

$$\Rightarrow (x-2400)(x+2000) = 0$$

$$\Rightarrow x-2400 = 0 \text{ or } x+2000 = 0$$

$$\Rightarrow x = 2400 \text{ or } -2000 \quad [1]$$

Since, the speed of aircraft cannot be negative
 So, $x = 2400$


Hence, the original speed is 2400 km/h.

\therefore Original duration of the flight

$$= \frac{6000}{2400} = 2 \frac{1}{2} \text{ hours} \quad \left[\because \text{time} = \frac{\text{distance}}{\text{speed}} \right] \quad [1]$$

36. In $\triangle ABC$, Given that F, E and D are the midpoints of AB, AC and BC respectively.

Hence, $FE \parallel BC$, $DE \parallel AB$ and $DF \parallel AC$



By mid-point theorem.

If $DE \parallel BA$

then $DE \parallel BF$

and if $FE \parallel BC$

then $FE \parallel BD$

\therefore FEDB is parallelogram in which DF is diagonal and a diagonal of Parallelogram divides it into two equal areas.

Hence $ar(\Delta BDF) = ar(\Delta DEF)$... (i) [½]

Similarly $ar(\Delta CDE) = ar(\Delta DEF)$... (ii) [½]

or $ar(\Delta AFE) = ar(\Delta DEF)$... (iii) [½]

or $ar(\Delta DEF) = ar(\Delta DEF)$... (iv) [½]

On adding eqns. (i), (ii), (iii) and (iv),


$$ar(\Delta BDF) + ar(\Delta CDE) + ar(\Delta AFE) + ar(\Delta DEF) \\ = 4ar(\Delta DEF)$$

or, $ar(\Delta ABC) = 4ar(\Delta DEF)$

$$\frac{ar(\Delta DEF)}{ar(\Delta ABC)} = \frac{1}{4} \quad [1]$$

OR

Given: A $\triangle ABC$ in which $AD \perp BC$ and $AD^2 = BD \cdot CD$.



To Prove: $\angle BAC = 90^\circ$.

$$\text{Proof: } AD^2 = BD \cdot CD \Rightarrow \frac{BD}{AD} = \frac{AD}{CD}$$

Now, in $\triangle DBA$ and $\triangle DAC$, we have:

$$\angle BDA = \angle ADC = 90^\circ \text{ and } \frac{BD}{AD} = \frac{AD}{CD}$$

$\therefore \triangle DBA \sim \triangle DAC$ [by SAS-similarity]

$\therefore \angle B = \angle 2$ and $\angle 1 = \angle C$


$$\therefore \angle 1 + \angle 2 = \angle B + \angle C \Rightarrow \angle A = \angle B + \angle C$$

$$\Rightarrow 2\angle A = \angle A + \angle B + \angle C = 180^\circ$$

$$\Rightarrow \angle A = 90^\circ.$$

37. Given: A circle of radius 4 cm.

Required: A pair of tangents, such that angle between them is 60° .



[2]

Step of Construction :

(i) Draw a circle of radius 4 cm with centre O .

(ii) Draw any radius OA .

(iii) Draw another radius OB , such that $\angle AOB$

$$= 120^\circ.$$

$$\therefore \angle P + \angle O = 180^\circ$$

$$\Rightarrow 60^\circ + \angle O = 180^\circ$$

$$\Rightarrow \angle O = 120^\circ$$

(iv) At point A, draw AP perpendicular to OA .

(v) At point B, draw BP perpendicular to OB and let the perpendiculars meet at P , such that $\angle P = 60^\circ$.

Thus, PA and PB are the required tangents. [2]

38. Let the first average speed of the bus be x km/h.

$$\therefore \frac{75}{x} + \frac{90}{x+10} = 3$$

$$\Rightarrow 75x + 750 + 90x = 3(x^2 + 10x) \quad [1]$$

$$\Rightarrow x^2 - 45x - 250 = 0 \quad [1]$$

$$\text{Solving to get } x = 50 \quad [1]$$

$$\therefore \text{Speed} = 50 \text{ km/h.} \quad [1]$$

CBSE Topper's Answer, 2015

Let the avg. speed for a dist. of 75 km = x km/hr

Then, time taken to cover 75 km = $\frac{75}{x}$ hrs

Now, speed for the next 90 km = $(x+10)$ km/hr

Time taken to cover 90 km = $\frac{90}{x+10}$ hrs.

A/Q

$$\frac{75}{x} + \frac{90}{x+10} = 3$$

$$\text{or, } 18 \left[\frac{5}{x} + \frac{6}{x+10} \right] = 3$$

$$\text{or, } \frac{5(x+10) + 6x}{x^2 + 10x} = \frac{1}{5}$$

$$\text{or, } 5x + 50 + 6x = \frac{x^2 + 10x}{5}$$

$$\text{or, } (11x + 50)5 = x^2 + 10x$$

$$\text{or, } x^2 + 10x - 55x - 250 = 0$$

$$\text{or, } x^2 - 45x - 250 = 0, \text{ which is a Quad eqn}$$

$$\text{or, } x^2 - 50x + 5x - 250 = 0$$

$$\text{or, } x(x-50) + 5(x-50) = 0$$

$$\text{or, } (x-50)(x+5) = 0$$

$$\text{or, } x-50 = 0 \quad | \quad \text{or, } x+5 = 0$$

$$\text{or, } x = 50 \quad | \quad \text{or, } x = -5$$

(invalid)

Ans: $\therefore \text{speed} = x = 50 \text{ km/hr}$

39. Volume of rain water on the roof = Volume of cylindrical tank [1]

i.e., $22 \times 20 \times h = \frac{22}{7} \times 1 \times 1 \times 3.5 \Rightarrow h = \frac{1}{40} \text{ m} = 2.5 \text{ cm}$ [2]

Water conservation must be encouraged or views relevant to it. [1]

CBSE Topper's Answer, 2015

radius of cylindrical tank = $\frac{2}{2} = 1 \text{ m}$.
 its height = $3.5 \text{ m.} = \frac{35}{10} \text{ m}$.
 Let the height of water on roof be h .
Volume of water on roof = Volume of water in tank.
 $\pi r^2 h = \pi r^2 h'$
 $22 \times 20 \times h = \frac{22}{7} \times \frac{22}{7} \times \frac{35}{10} \times 1 \times 1$
 $h = \frac{22}{2} \times \frac{1}{22} \times \frac{1}{20} \times \frac{35}{7} = \frac{1}{40} \text{ m}$
 $\therefore h = \frac{1}{40} \text{ m} = \frac{1}{40} \times 100 \text{ cm}$
 $= 2.5 \text{ cm}$
So, the rainfall is 2.5 cm

Views on water conservation:

- It is important practice in today's era of irrational water consumption and pollution. It should be practised at municipal level at all places.
- It can be done by many simple ways even at domestic level.
- Doing it is a sign of environmental consciousness.
- Some methods of water conservation are rooftop/surface water harvesting, building small earthen dams, etc.
- This conserved water helps refill underground water bodies and so, all must practise water conservation for sustainable development.

40. Find the median of the students and how can get the median graphically


| Age of student | C.I. | c.f. |
|----------------|-------|--------|
| Less than 6 | 4–6 | 2 |
| Less than 8 | 6–8 | 6 |
| Less than 10 | 8–10 | 12 |
| Less than 12 | 10–12 | 22 |
| Less than 14 | 12–14 | 42 |
| Less than 16 | 14–16 | 67 |
| Less than 18 | 16–18 | 76 |
| | | N = 76 |

[2]

$$\text{Median} = \frac{N}{2}\text{th term}$$

$$= \frac{76}{2} = 38\text{th term}$$

Median class = 12 – 14



[2]

Hence Medium = 13.6

OR

The class 35–40 has maximum frequency. So, it is the modal class.

∴ $x_k = 35, f_k = 50, f_{k-1} = 34, f_{k+1} = 42$ and $h = 5$.

$$\begin{aligned} \text{Mode, } M_0 &= x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\} \\ &= 35 + \left\{ 5 \times \frac{(50 - 34)}{(2 \times 50 - 34 - 42)} \right\} = 35 + \left\{ 5 \times \frac{16}{24} \right\} \\ &= \left\{ 35 + \frac{10}{3} \right\} = (5 + 3.33) = 38.33. \end{aligned}$$

Hence, mode = 38.33.

MATHEMATICS – BASIC

SOLUTIONS

SAMPLE QUESTION PAPER

CBSE Class X Examination

8

Section ‘A’

1. (c) 2. (a) 3. (b) 4. (a)
 5. (d) 6. (d) 7. (d) 8. (b)
 9. (b) 10. (b)

Fill in the blanks

11. $a = -\frac{3}{2}$

Explanation

The given equation is $5x^2 - \frac{7}{2}x + 2a = 0$

putting $x = \frac{1}{2}$

$$5\left(-\frac{1}{2}\right)^2 - \frac{7}{2}\left(-\frac{1}{2}\right) + 2a = 0$$

$$\frac{5}{4} + \frac{7}{4} + 2a = 0$$

$$a = -\frac{3}{2}$$

12. $x + 63$

Explanation

Here $a = x - 7$

$$d = (x - 2) - (x - 7) = 5$$


$$\therefore a_{15} = a + 14d$$

$$= x - 7 + 14(5)$$

$$= x + 63$$

13. $b = -1$

Explanation



Applying mid-point formula for Y coordinate

$$\frac{b+b+4}{2} = 1$$

$$2b + 4 = 2$$

$$2b = -2$$

$$b = -1$$

14. $QB = 3 \text{ cm}$

Explanation

$\Delta AOP \sim \Delta BOQ$ (AA)

$$\therefore \frac{OA}{AP} = \frac{BO}{QB}$$

$$\Rightarrow \frac{6}{4} = \frac{4.5}{QB}$$

$$\Rightarrow QB = 3 \text{ cm}$$

15. 2

Explanation

$$\frac{\sin 30^\circ}{\cos 60^\circ} + \frac{\tan 45^\circ}{\cot 45^\circ}$$

$$= \frac{1/2}{1/2} + \frac{1}{1} = 2$$

Answer the following.

16. $\sin (45^\circ + Q) - \cos (45^\circ - Q)$

$$= \cos [90 - (45^\circ - Q)] - \cos (45^\circ - Q) = 0$$

$$= \cos (45^\circ - Q) - \cos (45^\circ - Q)$$

$$= 0$$

17. Let one root be α , then other root will be $-\alpha$, then other root will be $-\alpha$.

Given equation is $3x^2 - px + 5 = 0$


Here $a = 3$, $b = -p$ and $c = 5$

Sum of roots $\alpha + \beta = -b/a = -(-p)/3$

$$\therefore \alpha + (-\alpha) = \frac{-(-p)}{3}$$

$$0 = \frac{P}{3}$$

$$\therefore P = 0$$



18. In quad. $AOBT$, $\angle A = \angle B = 90^\circ$
Also $\angle A + \angle O + \angle B + \angle T = 360^\circ$
 $90^\circ + 130^\circ + 90^\circ + \angle T = 360^\circ$
 $\therefore \angle T = 50^\circ$

19. In ΔPAO , $(OP)^2 = (12)^2 + (5)^2$
 $\therefore (OP)^2 = 169$
 $\therefore OP = 13$

Now in ΔPOB

$$\begin{aligned}(PB)^2 &= (13)^2 - (3)^2 \\ &= 169 - 9 \\ &= 160\end{aligned}$$

$$\therefore PB = \sqrt{160} = 4\sqrt{10} \text{ cm}$$

20. $PA = PB$ (tangents from an external point)
 $\therefore \angle PAB = \angle PBA = x$ (isosceles triangle)
 \Rightarrow In ΔABP
 $x + x + 60^\circ = 180^\circ$ (Angle sum property)
 $\Rightarrow x = 60^\circ$
 $\therefore \Delta ABP$ is an equilateral triangle.
 $\therefore AB = 5 \text{ cm}$

Section ‘B’

21. $366 \text{ days} = 52 \text{ weeks} + 2 \text{ days}$ [1]
2 days can be MT, TW, WT, TF, FS, SS, SM
 $= 7$
 $\Rightarrow P = \frac{2}{7}$ [1]

(CBSE Marking Scheme, 2012)

22. Total number of outcomes = 36
Favourable outcomes are (2, 6), (3, 5), (4, 4),
(5, 3), (6, 2) $= 5$ [1]
Required probability $= \frac{5}{36}$ [1]

(CBSE Marking Scheme, 2012)

23. $3 \times 12 \times 101 + 4 = 4(3 \times 3 \times 101 + 1)$
 $= 4(909 + 1)$ [1]
 $= 2 \times 2 \times 2 \times 5 \times 7 \times 13$
 $=$ a composite number [1]
[\because Product of more than two prime factors]
(CBSE Marking Scheme, 2015)

24. Given, $kx - 4y - 3 = 0$
and $6x - 12y - 9 = 0$
where, $a_1 = k$, $b_1 = -4$, $c_1 = -3$
 $a_2 = 6$, $b_2 = -12$, $c_2 = -9$
condition for infinite solution: $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ [1]
 $\frac{k}{6} = \frac{-4}{-12} = \frac{3}{9}$
Hence, $k = 2$ [1]

25. Let the first term be a and common difference d given
 $5a_5 = 8a_8$
 $5(a + 4d) = 8(a + 7d)$
 $5a + 20d = 8a + 56d$
 $3a + 36d = 0$ [1]
 $3(a + 12d) = 0$
 $a + 12d = 0$
 $a_{13} = 0$ [1]
(CBSE Marking Scheme, 2012)

26. Let the point on y -axis be $P(0, y)$
and $AP : PB = K : 1$
Therefore $\frac{5-K}{K+1} = 0$ gives $K = 5$
Hence required ratio is $5 : 1$ [1]

$$y = \frac{-4(5)-6}{6} = \frac{-13}{3}$$


Hence point on y -axis is $\left(0, \frac{-13}{3}\right)$. [1]

Section ‘C’

27. $P(x, y)$, $A(6, 2)$, $B(-2, 6)$
 $PA = PB$ or, $PA^2 = PB^2$
 $(x - 6)^2 + (y - 2)^2 = (x + 2)^2 + (y - 6)^2$ [1]
or $x^2 - 12x + 36 + y^2 - 4y + 4 = x^2 + 4x + 4$
 $+ y^2 - 12y + 36$
 $-12x - 4y = 4x - 12y$ [1]
 $12y - 4y = 4x + 12x$

$$\begin{aligned} 8y &= 16x \\ y &= 2x \end{aligned} \quad [1]$$

(CBSE Marking Scheme, 2015)

OR

$$\begin{aligned} \text{Mid point of BA} &= \frac{3+(-5)}{2} \\ &= -1 \text{ and } \frac{2-6}{2} = -2 \end{aligned}$$

$$D = (-1, -2)$$

$$\begin{aligned} \text{Mid points of BC} &= \frac{-5+7}{2} \\ &= 1 \text{ and } \frac{-6+4}{2} = -1 \end{aligned}$$

$$E = (1, -1)$$

Mid points of

$$CA = \frac{7+3}{2} = 5 \text{ and } \frac{4+2}{2} = 3$$

$$F = (5, 3)$$



Now, Area of ΔDEF

$$\begin{aligned} &= \frac{1}{2}[-1(-1-3)+1(3+2)+5(-2+1)] \\ &= \frac{1}{2}[4+5-5] \\ &= 2 \text{ units} \quad [1+1+1] \end{aligned}$$

(CBSE Marking Scheme, 2012)

- 28.** $AC = 8 \text{ cm}$, $AB = 10 \text{ cm}$ & $BC = 12 \text{ cm}$

$$\begin{aligned} \text{Let} \quad CF &= x \\ CF &= EC = x \\ AF &= 8-x = AD \\ BE &= 12-x = BD \quad [1] \end{aligned}$$

**29.**

$$\frac{AP}{AB} = \frac{3.5}{10.5} = \frac{1}{3} \quad [1]$$

$$\text{or} \quad \frac{AQ}{AC} = \frac{3}{9} = \frac{1}{3}$$

In ΔABC , $\frac{AP}{AB} = \frac{AQ}{AC}$ and $\angle A$ is commonor $\Delta APQ \sim \Delta ABC$

(By SAS) [1]


$$\frac{AP}{AB} = \frac{PQ}{BC}$$

$$\text{or} \quad \frac{1}{3} = \frac{4.5}{BC}$$

$$\text{or} \quad BC = 13.5 \text{ cm} \quad [1]$$

OR

$$\text{To prove: } \frac{ar(\Delta ABC)}{ar(\Delta DBC)} = \frac{AO}{DO}$$



[1]

Construction : Draw $AE \perp BC$ and $DF \perp BC$ **Proof :** In $\triangle AOE$ and $\triangle DOF$,

$$\angle AOE = \angle DOF$$

(vertically OPP. angles)

$$\angle AEO = \angle DFO = 90^\circ$$

(Construction)

$$\triangle AOE \sim \triangle DOF$$

(By AA similarity)

$$\therefore \frac{AO}{DO} = \frac{AE}{DF} \dots(i) \quad [1]$$

$$\text{Now, } \frac{ar(\Delta ABC)}{ar(\Delta DBC)} = \frac{\frac{1}{2} \times BC \times AE}{\frac{1}{2} \times BC \times DF}$$

$$= \frac{AE}{DF}$$

$$= \frac{AO}{DO} \quad [\text{From eq. (i)}] \quad [1]$$

Hence proved

30.

$$\begin{array}{r} x^2 + 2x \\ \overline{x^3 + 5x^2 + 7x + 3} \\ - - \\ 3x^2 + 7x + 3 \\ - - \\ 3x^2 - 6x \\ - - \\ x + 3 \end{array} \quad [2]$$

Remainder = $x + 3$

Hence, $-(x + 3)$ must be added [1]

(CBSE Marking Scheme, 2016)

31. Let $3 + \sqrt{5}$ be a rational number.

$$\therefore 3 + \sqrt{5} = \frac{p}{q}, q \neq 0 \quad [1]$$

$$3 + \sqrt{5} = \frac{p}{q}$$

$$\Rightarrow \sqrt{5} = \frac{p-3}{q}$$

$$\Rightarrow \sqrt{5} = \frac{p-3q}{q}$$

$\sqrt{5}$ is irrational and $\frac{p-3q}{q}$ is rational [1]

But rational number can not be equal to an irrational number.

$\therefore 3 + \sqrt{5}$ is an irrational number. [1]

32. We have:

$$\begin{aligned} \text{LHS} &= \frac{(1 - \cos \theta)}{(1 + \cos \theta)} \\ &= \frac{(1 - \cos \theta)}{(1 + \cos \theta)} \times \frac{(1 - \cos \theta)}{(1 - \cos \theta)} \quad [1] \\ &[\text{multiplying num. and denon. by } (1 - \cos \theta)] \\ &= \frac{(1 - \cos \theta)^2}{(1 - \cos^2 \theta)} \quad [1] \\ &= \frac{(1 + \cos^2 \theta - 2\cos \theta)}{\sin^2 \theta} \\ &= \left(\frac{1}{\sin^2 \theta} + \frac{\cos^2 \theta}{\sin^2 \theta} - \frac{2\cos \theta}{\sin^2 \theta} \right) \\ &= (\cosec^2 \theta + \cot^2 \theta - 2\cosec \theta \cot \theta) \\ &\quad [1] \\ &= (\cosec \theta - \cot \theta)^2 = \text{RHS.} \end{aligned}$$

LHS = RHS

OR

$$\begin{aligned} \text{LHS} &= \frac{\cos A}{1 + \tan A} - \frac{\sin A}{1 + \cot A} \\ &= \frac{\cos A}{1 + \frac{\sin A}{\cos A}} - \frac{\sin A}{1 + \frac{\cos A}{\sin A}} \\ &= \frac{\cos^2 A}{\cos A + \sin A} - \frac{\sin^2 A}{\sin A + \cos A} \\ &= \frac{\cos^2 A - \sin^2 A}{(\sin A + \cos A)} \\ &= \frac{(\cos A + \sin A)(\cos A - \sin A)}{\sin A + \cos A} \\ &= \cos A - \sin A \\ &= \text{RHS} \quad \text{Hence Proved} \end{aligned}$$

33. Volume of water in cylinder

$$\begin{aligned} &= \text{volume of cylinder} \\ &= \pi r^2 h \\ &= \pi \times (60)^2 \times 180 \\ &= 648000\pi \text{ cm}^3 \end{aligned}$$

water displaced on dropping cone, volume of

$$\begin{aligned} \text{solid cone} &= \frac{1}{3} \pi r^2 h \\ &= \frac{1}{3} \times \pi \times (30)^2 \times 60 \\ &= 18000\pi \text{ cm}^3 \end{aligned}$$

Volume of water left in cylinder = volume of cylinder – volume of cone

$$\begin{aligned} &= 648000\pi - 18000\pi \\ &= 630000\pi \text{ cm}^3 \end{aligned}$$

$$\begin{aligned} &= \frac{630000 \times 22}{1000000 \times 7} \text{ m}^3 \\ &= 1.98 \text{ m}^3 \end{aligned}$$

[3] (CBSE Marking Scheme, 2015)

OR

Required area

= (area of rect. ABCD) – (area of semicircle with $r = 7 \text{ cm}$)

$$= \left[(18 \times 14) - \left(\frac{1}{2} \times \pi \times r^2 \right) \right] \text{ cm}^2$$

$$= \left(252 - \frac{1}{2} \times \frac{22}{7} \times 7 \times 7 \right) \text{ cm}^2$$

$$= (252 - 77) \text{ cm}^2 = 175 \text{ cm}^2.$$

34.

| Classes | Frequency f_i | Mid-point x_i | $f_i x_i$ |
|---------|-----------------|-----------------|-----------------------|
| 0–20 | 6 | 10 | 60 |
| 20–40 | 8 | 30 | 240 |
| 40–60 | 10 | 50 | 500 |
| 60–80 | 12 | 70 | 840 |
| 80–100 | 8 | 90 | 720 |
| 100–120 | 6 | 110 | 660 |
| | $\sum f_i = 50$ | | $\sum f_i x_i = 3020$ |

[2]

$$\begin{aligned} \text{Mean } \bar{x} &= \frac{\sum f_i x_i}{\sum f_i} = \frac{3020}{50} \\ &= 60.4 \end{aligned}$$

[1]

Section 'D'

35. Let the speed while going be $x \text{ km/h}$

\therefore Speed while returning = $(x + 10) \text{ km/h}$

According to question,

$$\frac{150}{x} - \frac{150}{x+10} = \frac{5}{2} \quad [1]$$

$$\Rightarrow \frac{150(x+10) - x(150)}{x(x+10)} = \frac{5}{2}$$

$$\Rightarrow \frac{150x + 1500 - 150x}{x^2 + 10x} = \frac{5}{2}$$

$$\begin{aligned} \Rightarrow 5(x^2 + 10x) &= 2 \times 1500 \\ \Rightarrow x^2 + 10x &= 600 \quad [1] \end{aligned}$$

$$\Rightarrow x^2 + 10x - 600 = 0$$

$$\Rightarrow (x + 30)(x - 20) = 0$$

$$\text{or } x = 20$$

$$\therefore \text{Speed while going} = 20 \text{ km/h} \quad [1]$$

$$\begin{aligned} \text{and } \text{speed while returning} &= 20 + 10 \\ &= 30 \text{ km/h} \quad [1] \end{aligned}$$

[CBSE Marking Scheme, 2016]

OR

Here $x = -2$ is the root of the equation

$$3x^2 + 7x + P = 0$$

$$\text{or } 3(-2)^2 + 7(-2) + P = 0$$

$$\Rightarrow P = 2$$


Root of the equation $x^2 + 4kx + k^2 - k + 2 = 0$ are equal.

or $16k^2 - 4(k^2 - k + 2) = 0$
 $16k^2 - 4k^2 + 4k - 8 = 0$
 $12k^2 + 4k - 8 = 0$
 $3k^2 + k - 2 = 0$
 $(3k - 2)(k + 1) = 0$
 $k = \frac{2}{3}, -1$

Hence, roots = $\frac{2}{3}, -1$

[4] [CBSE Marking Scheme, 2015]

36.



[1]

Let $BD = DE = EC$ be x
 and $BE = 2x$
 $BC = 3x$ [1]

Now in $\triangle ABE$, $AE^2 = AB^2 + BE^2$
 $= AB^2 + 4x^2$... (i)

In $\triangle ABC$, $AC^2 = AB^2 + BC^2 = AB^2 + 9x^2$

In $\triangle ADB$, $AD^2 = AB^2 + BD^2 = AB^2 + x^2$

Now, on multiplying (i) by 8

$$8 AE^2 = 8 AB^2 + 32x^2 \quad \dots (ii) [1]$$

$$3 AC^2 + 5 AD^2 = 3 (AB^2 + 9x^2) + 5 (AB^2 + x^2)$$

$$= 3 AB^2 + 27x^2 + 5 AB^2 + 5x^2$$

$$= 8 AB^2 + 32x^2 = 8$$

$$AB^2 + 4x^2 \quad \dots (iii)$$


$$\therefore 3 AC^2 + 5 AD^2 = 8 AE^2 \quad [1]$$

[From eq. (ii) & (iii)]

Hence Proved

OR

Given: A $\triangle ABC$ in which $\angle C = 90^\circ$. P and Q are points on CA and CB respectively.



To Prove: $(AQ^2 + BP^2) = (AB^2 + PQ^2)$.

Proof: From right $\triangle ACQ$, we have:

$$AQ^2 = (AC^2 + CQ^2) \quad \dots (i)$$

[by Pythagoras' theorem]

From right $\triangle BCP$, we have:

$$BP^2 = (BC^2 + CP^2) \quad \dots (ii)$$

[by Pythagoras' theorem]

From right $\triangle ACB$, we have:

$$AB^2 = AC^2 + BC^2 \quad \dots (iii)$$

[by Pythagoras' theorem]

From right $\triangle PCQ$, we have:

$$PQ^2 = (CQ^2 + CP^2) \quad \dots (iv)$$

[by Pythagoras' theorem]

From (i) and (ii), we get:

$$(AQ^2 + BP^2) = (AC^2 + BC^2) + (CQ^2 + CP^2) = (AB^2 + PQ^2) \quad [\text{using (iii) and (iv)}].$$

Hence, $(AQ^2 + BP^2) = (AB^2 + PQ^2)$


37. Steps of Construction:

(i) Draw a line segment $AB = 5\text{cm}$

(ii) At A make $\angle BAY = 45^\circ$

(iii) Take A as centre and radius $AC = 6\text{ cm}$, draw an arc cutting AY at C. [1]

(iv) Join BC to obtain the $\triangle ABC$.



(v) Draw any ray AX making an acute angle with AB on the side opposite to the vertex C. [1]

(vi) Locate 6 points (the greater of 6 and 5 in A_1, A_2, A_3, A_4, A_5 and A_6) on AX, such that


$$AA_1 = A_1A_2 = A_2A_3 = A_3A_4 = A_4A_5 = A_5A_6 \quad [1]$$

(vii) Join the 5th point, being the smaller of 5 and 6 in 6/5 to B and draw a line segment through A_6 parallel A_5B intersecting the extended line segment AB at B' .

(viii) Draw a line through B parallel to BC intersecting the extended line segment AC at C' .

Then $AB'C'$ is the required triangle. [1]

38. Let AB is a building 60 m high and is a tower h meter high. Angle of depressions of top and bottom are given 30° and 60° respectively



$$DC = EB = h \text{ m and}$$

Let $BC = x$

$$\Rightarrow AE = (60 - h)\text{m}$$

$$\text{In } \triangle AED, \frac{60-h}{ED} = \tan 30^\circ \quad [1\frac{1}{2}]$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{60-h}{BC}$$

$$\Rightarrow \sqrt{3}(60-h) = x \quad \dots(i) [1\frac{1}{2}]$$

$$\text{In } \triangle ABC, \frac{60}{x} = \tan 60^\circ$$

$$60 = \sqrt{3}x \quad \dots(ii) [1]$$

Putting the value of x from equation (i) in equation (ii), we get

$$60 = \sqrt{3} \times \sqrt{3}(60-h)$$

$$60 = 3 \times (60-h)$$

$$20 = 60 - h$$

$$h = 40 \text{ m}$$

Hence, Height of tower = 40m. [1]

39. Inner radius of the glass = $\frac{7}{2}$ cm and height = 16 cm.

Apparent capacity of the glass = $\pi r^2 h$

$$= \left(\frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times 16 \right) \text{cm}^3 \\ = 616 \text{ cm}^3. \quad [1\frac{1}{2}]$$

Volume of the hemisphere in the bottom

$$= \frac{2}{3} \pi r^3 = \left(\frac{2}{3} \times \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times \frac{7}{2} \right) \text{cm}^3 \\ = \frac{539}{6} \text{ cm}^3 = 89.83 \text{ cm}^3. \quad [1\frac{1}{2}]$$

Actual capacity of the glass

$$= (\text{volume of the glass}) - (\text{volume of the hemisphere}) \\ = (616 - 89.83) \text{ cm}^3 = 526.17 \text{ cm}^3. \quad [1]$$

| Classes | f | c.f. |
|---------|-----|------|
| 5–10 | 2 | 2 |
| 10–15 | 12 | 14 |
| 15–20 | 2 | 16 |
| 20–25 | 4 | 20 |
| 25–30 | 3 | 23 |
| 30–35 | 4 | 27 |
| 35–40 | 3 | 30 |

[2]

$$\text{Total} \quad \sum f = 30 = N$$

$$\text{Since,} \quad \text{Median} = \frac{N}{2} \text{th term}$$

$$= \frac{30}{2} = 15 \text{th term} \quad [1\frac{1}{2}]$$

Median class = 15–20

[½]

$$\text{Medain} = l + \left(\frac{\frac{N}{2} - c.f.}{f} \right) \times h$$


$$l = 15, N = 30, c.f. = 14, f = 2 \text{ & } h = 5$$

$$\begin{aligned} \text{Median} &= 15 + \left(\frac{15-14}{2} \right) \times 5 \\ &= 15 + 2.5 = 17.5 \end{aligned}$$

[1]

OR

| x | y |
|--------------|-----|
| More than 10 | 100 |
| More than 20 | 90 |
| More than 30 | 75 |
| More than 40 | 45 |
| More than 50 | 13 |
| More than 60 | 5 |
| More than 70 | 0 |



MATHEMATICS – BASIC

SOLUTIONS SAMPLE QUESTION PAPER

CBSE Class X Examination

9

Section 'A'

[1 × 20 = 20]

1. (a) 2. (b) 3. (b) 4. (a)
5. (b) 6. (a) 7. (b) 8. (b)
9. (d) 10. (b)

Fill in the blanks

11. $k = 3$

Explanation

Let one zero be α , then other will be $\frac{1}{\alpha}$

Given equation is $3x^2 + 8x + k = 0$

Here $a = 3, b = 8$ and $c = k$

$$\text{product of zeroes} = \alpha \cdot \frac{1}{\alpha} = \frac{c}{a}$$

$$\Rightarrow 1 = \frac{k}{3}$$

$$\Rightarrow k = 3$$

12. $-2/3$

Explanation

Given equation is $kx^2 + 2x + 3k = 0$

Here $a = k, b = 2$ and $c = 3k$

Sum of zeroes = product of zeroes

$$\frac{-b}{a} = \frac{c}{a}$$

$$\frac{-2}{k} = \frac{3k}{k} \quad \text{or} \quad k = -2/3$$

13. $a = 3$

Explanation

Applying mid-point formula for Y -coordinate

$$\frac{7+a}{2} = 5$$

$$\Rightarrow a = 3$$

14. 10/9

Explanation

Given equation is

$$x^2 - 2(1 + 3k)x + 7(3 + 2k) = 0$$

Here $a = 1, b = -2(1 + 3k)$ and $c = 7(3 + 2k)$

For equal roots,

$$b^2 - 4ac = 0$$

$$[-2(1 + 3k)]^2 - 4 \times 1 \times 7(3 + 2k) = 0$$

$$4(1 + 3k)^2 - 4 \times (21 + 14k) = 0$$

$$4[1 + 9k^2 + 6k - 21 - 14k] = 0$$

$$9k^2 - 8k - 20 = 0$$

$$9k^2 - 18k + 10k - 20 = 0$$

$$9k(k - 2) + 10(k - 2) = 0$$

$$(k - 2)(9k + 10) = 0$$

$$\Rightarrow k = 2 \quad \text{or} \quad 10/9$$

15. 24

Explanation

Here $a = 3$ and $d = 7$

$$\begin{aligned} \therefore a_4 &= a + 3d \\ &= 3 + 3 \times 7 \\ &= 24 \end{aligned}$$

Answer the following.

16. $(k + 1), 3k, (4k + 2)$

Common difference

$$3k - (k + 1) = (4k + 2) - 3k$$

$$2k - 1 = k + 2$$

$$k = 3$$

17.
$$\sqrt{\frac{\sec A - \tan A}{\sec A + \tan A} \times \frac{\sec A - \tan A}{\sec A - \tan A}}$$

$$= \sqrt{\frac{(\sec A - \tan A)^2}{\sec^2 A - \tan^2 A}}$$


$$= \sqrt{\frac{(\sec A - \tan A)^2}{1}} \quad (\sec^2 A - \tan^2 A = 1)$$

$$= \sec A - \tan A$$

$$\begin{aligned} 18. &= [(\sin^2 \theta)^2 - (\cos^2 \theta)^2 + 1] \operatorname{cosec}^2 \theta \\ &= [(\sin^2 \theta + \cos^2 \theta) (\sin^2 \theta - \cos^2 \theta) + 1] \operatorname{cosec}^2 \theta \\ &= [\sin^2 \theta - \cos^2 \theta + 1] \operatorname{cosec}^2 \theta \\ &= [\sin^2 \theta + (1 - \cos^2 \theta)] \operatorname{cosec}^2 \theta \\ &= [\sin^2 \theta + \sin^2 \theta] \operatorname{cosec}^2 \theta \\ &= 2 \sin^2 \theta \times \frac{1}{\sin^2 \theta} = 2 \end{aligned}$$

19. Area of triangle formed by three points is zero if they are collinear

$$\begin{aligned} &\Rightarrow \frac{1}{2} [2(k-7) + 5(7-3) + 6(3-k)] = 0 \\ &\Rightarrow 2k - 14 + 20 + 18 - 6k = 0 \\ &\quad -4k + 24 = 0 \\ &\quad k = 6 \end{aligned}$$



$$OB^2 = (24)^2 + (10)^2$$

$$= 576 + 100$$

$$OB^2 = 676 = (26)^2$$

$$OB = 26$$

Section ‘B’

21. Let the point p lie on x-axis

\therefore P (x, 0) is equidistant from point A (2, -5) and B (-2, 9)

$$\therefore PA^2 = PB^2$$

$$\text{or}, (2-x)^2 + (5-0)^2$$

$$= (-2-x)^2 + (9-0)^2$$

[1]

$$\begin{aligned} &\Rightarrow 4 - 4x + x^2 + 25 = 4 + 4x + x^2 + 81 \\ &\Rightarrow -8x = 56 \\ &\Rightarrow x = -7 \end{aligned} \quad [1]$$

\therefore The point is (-7, 0).

22. Here, $a_1 = -1, a_2 = -5$ and $d = 4$

$$S_n = \frac{n}{2} [2a + (n-1)d] \quad [1]$$

$$\therefore S_{16} = \frac{16}{2} [2 \times (-1) + (16-1)(-4)]$$

$$S_{16} = 8 [-2 - 60] = 8 (-62)$$

$$\therefore S_{16} = -496 \quad [1]$$

23. Here, $a_1 = 2, b_1 = 1, c_1 = -3$

and $a_2 = 4, b_2 = 2, c_2 = -6$

If $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ then the lines are coincident

$$\text{clearly } \frac{2}{4} = \frac{1}{2} = \frac{3}{6} \quad [1]$$

Hence lines are coincident. [1]

24. $240 = 228 \times 1 + 12$

$$228 = 12 \times 19 + 0$$

Hence, HCF of 240 and 228 = 12. [2]

25. No. of cards = 20

Multiples of 5 from 11 to 30 are 15, 20, 25 and 30 so, number of favourable outcomes = 4 [1]

$$\therefore \text{Required probability} = \frac{4}{20} = \frac{1}{5} \quad [1]$$

26. When two dices are thrown all possible outcomes = $6 \times 6 = 36$

If sum of both faces should be 10, they are $\{(4, 6), (6, 4), (5, 5)\}$

\therefore No. of favourable outcomes = 3 [1]

$$\therefore \text{Required probability} = \frac{3}{36} = \frac{1}{12} \quad [1]$$

Section ‘C’

$$27. x^2 - 2\sqrt{2}x = 0$$

$$x(x - 2\sqrt{2}) = 0$$

Zeroes are 0 and $2\sqrt{2}$. [1]

$$\text{Sum of zeroes} = 2\sqrt{2} = -\frac{\text{coefficient of } x}{\text{coefficient of } x^2}$$

$$\text{and product of zeroes} = \frac{2\sqrt{2}}{1} \quad [1]$$

$$0 = \frac{\text{constant term}}{\text{coefficient of } x^2} = 0. \quad [1]$$

28. $92 = 2^2 \times 23$

$$510 = 2 \times 3 \times 5 \times 17$$

$$\text{HCF}(510, 92) = 2 \quad [1]$$

$$\begin{aligned} \text{LCM}(510, 92) &= 2^2 \times 23 \times 3 \times 5 \times 17 \\ &= 23460 \end{aligned} \quad [1]$$

$$\begin{aligned} \text{HCF}(510, 92) \times \text{LCM}(510, 92) \\ &= 2 \times 23460 = 46920 \end{aligned}$$

Product of two number

$$= 510 \times 92 = 46920$$

Hence, HCF \times LCM

= Product of two numbers.

[1]

29. We know that $AB \parallel DC$ in trap. ABCD and its diagonals intersect at O. Then, we have:

$$\frac{AO}{OC} = \frac{BO}{OD} \Rightarrow \frac{3x-1}{5x-3} = \frac{2x+1}{6x-5} \quad [1]$$

$$\Rightarrow (3x-1)(6x-5) = (2x+1)(5x-3)$$

$$\Rightarrow 18x^2 - 21x + 5 = 10x^2 - x - 3$$

$$\Rightarrow 8x^2 - 20x + 8 = 0$$

$$\Rightarrow 2x^2 - 5x + 2 = 0$$

$$\Rightarrow (x-2)(2x-1) = 0$$

$$\Rightarrow x = 2 \text{ or } x = \frac{1}{2}. \quad [1]$$

But, $x = \frac{1}{2}$ will make $OC = (5x-3) \text{ cm}$


$$= \left(5 \times \frac{1}{2} - 3\right) \text{ cm} = -\frac{1}{2} \text{ cm.} \quad [1]$$

And, the distance cannot be negative.

$$\therefore x \neq \frac{1}{2}.$$

Hence, $x = 2$.

OR



Given,

$$\begin{aligned} DE &\parallel BC \\ \frac{AD}{DB} &= \frac{AE}{EC} \end{aligned} \quad (\text{By BPT})$$

$$\text{or, } \frac{x+2}{3x+16} = \frac{x}{3x+5}$$

$$(x+2)(3x+5) = x(3x+16)$$

$$3x^2 + 5x + 6x + 10 = 3x^2 + 16x$$


$$11x + 10 = 16x$$

$$16x - 11x = 10$$

$$5x = 10$$

$$x = 2$$

30.



Since the tangent is perpendicular to the end point of radius,

$$\begin{aligned} \angle OBP &= 90^\circ \\ \angle OAB &= \angle OBA \quad (\because OA = OB) \\ \angle OBA &= 30^\circ \end{aligned} \quad [1]$$


$$\therefore \angle AOB = 180^\circ - (30^\circ + 30^\circ)$$

$$\angle AOB = 120^\circ$$

$$\angle ABP = \angle OBP - \angle OBA$$

$$\angle ABP = 90^\circ - 30^\circ = 60^\circ \quad [1]$$

31.



$$AP = \frac{2}{5} AB$$

$$\begin{aligned}
 \text{or, } AP : PB &= 2 : 3 & [1] & \therefore \text{Area of triangle} \\
 x &= \frac{2 \times 6 + 3 \times 1}{2+3} & [\frac{1}{2}] & \frac{1}{2} [x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2)] & [\frac{1}{2}] \\
 \text{and } y &= \frac{2 \times 7 + 3 \times 2}{2+3} & [\frac{1}{2}] & \text{or, } \frac{1}{2} [5(4-y) + (-3)(y-2) + x(2-4)] = 0 \\
 \therefore x &= \frac{12+3}{5} = 3; & & \frac{1}{2} [20 - 5y - 3y + 6 + (-2x)] = 0 \\
 y &= \frac{14+6}{5} = 4 & & \frac{1}{2} [-2x - 8y + 26] = 0 \\
 P(x, y) &= (3, 4) & [1] & x + 4y - 13 = 0 \\
 \text{OR} & & & & [1] \\
 \text{Since the points are collinear} & & & & \\
 \therefore \text{The area of triangle} &= 0 & & & \text{Hence Proved}
 \end{aligned}$$

32.

| Class | x_i (class marks) | f_i | $f_i x_i$ | c.f. |
|-------|---------------------|------------------|-----------------------|------|
| 0–10 | 5 | 8 | 40 | 8 |
| 10–20 | 15 | 16 | 240 | 24 |
| 20–30 | 25 | 36 | 900 | 60 |
| 30–40 | 35 | 34 | 1190 | 94 |
| 40–50 | 45 | 6 | 270 | 100 |
| | | $\sum f_i = 100$ | $\sum f_i x_i = 2640$ | |

[1]

$$\text{Mean} = \frac{\sum f_i x_i}{\sum f_i} = \frac{2640}{100} = 26.4 \quad R^3 = 8 \quad R = 2 \text{ cm} \quad [1]$$

$$\text{Median} = \frac{N}{2} \text{th term} \quad \text{OR} \quad \text{Side of the square} = \sqrt{121} \text{ cm} = 11 \text{ cm}$$

$$= \frac{100}{2} = 50 \text{th term} \quad \text{Perimeter of the square} = (4 \times 11) \text{ cm} = 44 \text{ cm.}$$

$$\text{Median Class} = 20–30 \quad \therefore \text{length of the wire} = 44 \text{ cm.}$$

$$\text{Median} = l_1 + \frac{\frac{N}{2} - cf}{f} \times h \quad \therefore \text{circumference of the circle} = \text{length of the wire} = 44 \text{ cm}$$

$$\text{Median} = 20 + \frac{50 - 24}{36} \times 10 \quad \text{Let the radius of the circle be } r \text{ cm.}$$

$$\text{Median} = 20 + 7.22 = 27.22 \quad \text{Then, } 2\pi r = 44 \Rightarrow 2 \times \frac{22}{7} r = 44 \Rightarrow r = 7.$$

$$33. \text{ Let the radius of sphere} = R \text{ cm} \quad \therefore \text{area of the circle} = \pi r^2$$

$$\text{Volume of sphere} = \text{Volume of cone} \quad = \left(\frac{22}{7} \times 7 \times 7 \right) \text{ cm}^2$$

$$\Rightarrow \frac{4}{3}\pi R^3 = \frac{1}{3}\pi r^2 h \quad [1] \quad = 154 \text{ cm}^2.$$

$$\Rightarrow \frac{4}{3}\pi R^3 = \frac{1}{3}\pi \times 2 \times 2 \times 8 \quad 34. \text{ According to the question,}$$

$$R^3 = \frac{2 \times 2 \times 8}{4} \quad \sin 3\theta = \cos(\theta - 6^\circ) \quad [1]$$

$$\cos(90^\circ - 3\theta) = \cos(\theta - 6^\circ) \quad [1]$$

$$90^\circ - 3\theta = \theta - 6^\circ \quad [1]$$

$$4\theta = 90^\circ + 6^\circ = 96^\circ \quad [1]$$

$$\theta = \frac{96^\circ}{4} = 24^\circ \quad [1]$$

OR

We have:

$$\begin{aligned}
 x^2 + y^2 &= (a \sin \theta + b \cos \theta)^2 + (a \cos \theta + b \sin \theta)^2 \\
 &= (a^2 \sin^2 \theta + b^2 \cos^2 \theta + 2ab \sin \theta \cos \theta) \\
 &\quad + (a^2 \cos^2 \theta + b^2 \sin^2 \theta - 2ab \sin \theta \cos \theta) \\
 &= a^2 \sin^2 \theta + b^2 \cos^2 \theta + a^2 \cos^2 \theta + b^2 \sin^2 \theta \\
 &= a^2(\sin^2 \theta + \cos^2 \theta) + b^2(\sin^2 \theta + \cos^2 \theta) \\
 &= a^2 + b^2 \quad [\because \sin^2 \theta + \cos^2 \theta = 1].
 \end{aligned}$$

Hence, $x^2 + y^2 = a^2 + b^2$.**Section 'D'**

35. Let the speed of the car I from A be
- x
- km/hr.

Speed of the car II from B be y km/hr

Same direction :

Distance covered by car I = $150 + (\text{distance covered by car II})$

$$\Rightarrow 15x = 150 + 15y$$

$$\Rightarrow 15x - 15y = 150$$

$$\Rightarrow x - y = 10 \quad \dots(i) [1]$$

Opposite Direction :

Distance covered by car I + distance covered by car II

$$= 150 \text{ km}$$

$$x + y = 150 \quad \dots(ii) [1]$$

Adding eq(i) and (ii),

$$2x = 160$$

$$\therefore x = 80$$

Substituting $x = 80$ in eq (i),

$$y = 70$$

 \therefore Speed of the car I from A = 80 km/hr [1]

and speed of the car II from B = 70 km/hr [1]

OR $x = -4$ is the root of the equation

$$x^2 + 2x + 4p = 0$$

$$(-4)^2 + 2(-4) + 4p = 0$$

$$16 - 8 + 4p = 0$$

$$p = -2$$

Equation $x^2 - 2(1+3k)x + 7(3+2k) = 0$ has equal roots

$$\therefore 4(1+3k)^2 - 28(3+2k) = 0$$


$$9k^2 - 8k - 20 = 0$$

$$(9k+10)(k-2) = 0$$

$$k = \frac{-10}{9}, 2$$

Hence, the value of $k = \frac{-10}{9}, 2$.

36. Let AB be the pole and let AC be its shadow.



[1]

Let the angle of elevation of the sun be θ° .Then, $\angle ACB = \theta$, $\angle CAB = 90^\circ$,

$$AB = 10 \text{ cm} \text{ and } AC = 10\sqrt{3} \text{ m.} \quad [1]$$

From right $\triangle CAB$, we have


$$\tan \theta = \frac{AB}{AC} = \frac{10}{10\sqrt{3}} = \frac{1}{\sqrt{3}} \quad [1]$$

$$\Rightarrow \theta = 30^\circ, \quad [1]$$

Hence, the angular elevation of the sun is 30° .

- 37.
- Steps of construction:**

- (i) Draw a line segment BC = 8 cm
- (ii) Draw perpendicular bisector of BC which intersects BC at O.



- (iii) Mark A on bisector such that $AO = 4 \text{ cm}$
(iv) Join A to B and A to C
(v) Draw an acute angle CBX at B fo BC, down word.
(vi) Mark B_1, B_2, B_3 on BX, such that $BB_1 = B_1B_2 = B_2B_3$
(vii)Join B_3 to C.
(viii)Draw $B_2C^1 \parallel B_3C$ to meet BC at C^1
(ix) From C^1 draw $C^1A^1 \parallel CA$, to meet AB at A^1 .
Hence A^1BC^1 is the required triangle.

[2 + 2] (CBSE Marking Scheme, 2017)

38. Volume of lead = $(44 \times 44 \times 44) \text{ cm}^3$.

Radius of one bullet = 2 cm.

Volume of one bullet

$$= \left(\frac{4}{3} \times \frac{22}{7} \times 2 \times 2 \times 2 \right) \text{cm}^3 \quad \left[\because V = \frac{4}{3} \pi r^3 \right]$$

$$= \left(\frac{704}{21} \right) \text{cm}^3. \quad [2]$$

Number of bullets made = $\frac{\text{volume of cube}}{\text{volume of 1 bullet}}$

$$= \frac{44 \times 44 \times 44}{\left(\frac{704}{21} \right)} = \left(44 \times 44 \times 44 \times \frac{21}{704} \right)$$


$$= 2541. \quad [2]$$

39. Given : DEF is a triangle in which $XY \parallel EF$

$$\text{To Prove : } \frac{DX}{XE} = \frac{DY}{YF}$$

Construction : Draw $XN \perp DY$ and $YM \perp DX$, Join EY and FX.

Proof : In $\triangle DEF$,



$$\text{area of } \triangle DXY = \frac{1}{2} \times DX \times YM \quad \dots(i)$$

$$\text{area of } \triangle XYE = \frac{1}{2} \times XE \times YM \quad \dots(ii)$$

Dividing eq (i) by eq (ii)

$$\frac{\text{area of } \triangle DXY}{\text{area of } \triangle XYE} = \frac{\left(\frac{1}{2} \right) \times (DX) \times (YM)}{\left(\frac{1}{2} \right) \times (XE) \times (YM)} \quad [1]$$

$$\frac{\text{area of } \triangle DXY}{\text{area of } \triangle XYE} = \frac{DX}{XE} \quad \dots(iii)$$

$$\text{area of } \triangle DXY = \left(\frac{1}{2} \right) \times (DY) \times (XN) \quad \dots(iv)$$

$$\text{area of } \triangle XYF = \left(\frac{1}{2} \right) \times (YF) \times (XN) \quad \dots(v)$$

Dividing eq (iv) by eq (v),

$$\frac{\text{area of } \triangle DXY}{\text{area of } \triangle XYF} = \frac{\left(\frac{1}{2} \right) \times (DY) \times (XN)}{\left(\frac{1}{2} \right) \times (YF) \times (XN)} \quad [1]$$

$$\frac{\text{area of } \triangle DXY}{\text{area of } \triangle XYF} = \frac{DY}{YF} \quad \dots(vi) [1]$$

$\triangle XYE$ and $\triangle XYF$ lie on the same base and between same parallel lines XY and EF.

$$\text{area } \triangle XYE = \text{area } \triangle XYF \quad \dots(vii)$$

From eq (vii),

$$\frac{\text{area of } \triangle DXY}{\text{area of } \triangle XYE} = \frac{DY}{YF} \quad \dots(viii)$$

On comparing eq (iv) and eq (viii)

$$\frac{DX}{XE} = \frac{DY}{YF} \quad [1]$$

Hence Proved

OR


In $\triangle ABC$ and $\triangle ADE$

$$\angle ABC = \angle ADE \quad [\text{Given}]$$

$$\angle A = \angle A \quad [\text{Common}]$$

$$\triangle ABC \sim \triangle ADE \quad [\text{AA Similarity}]$$

$$\frac{AB}{AD} = \frac{BC}{DE} = \frac{AC}{AE} \quad [1]$$



$$\begin{aligned}\frac{AB}{AD} &= \frac{BC}{DE} \\ \frac{AE+BE}{AD} &= \frac{BC}{DE} \\ \frac{7.2+4.2}{7.6} &= \frac{8.4}{DE} \\ \frac{11.4}{7.6} &= \frac{8.4}{DE} \\ DE &= \frac{8.4 \times 7.6}{11.4} \\ DE &= 5.6 \text{ cm}\end{aligned}$$

| Life Times | c.f. |
|----------------|------|
| Less than 1200 | 15 |
| Less than 1400 | 75 |
| Less than 1600 | 143 |
| Less than 1800 | 229 |
| Less than 2000 | 304 |
| Less than 2200 | 365 |
| Less than 2400 | 410 |

[1]

OR


| Class | x_i (class marks) | f_i | $f_i x_i$ |
|--------------|---------------------|-----------------|-------------------------|
| 0–100 | 50 | 12 | 600 |
| 100–200 | 150 | 16 | 2400 |
| 200–300 | 250 | 6 | 1500 |
| 300–400 | 350 | 7 | 2450 |
| 400–500 | 450 | 9 | 4050 |
| Total | | $\sum f_i = 50$ | $\sum f_i x_i = 11,000$ |

[2]

$$\text{Mean} = \frac{\sum f_i x_i}{\sum f_i} = \frac{11000}{50} = 220 \quad [1]$$

Average daily income = ₹ 220. [1]

[1]



[2]

[2]

MATHEMATICS – BASIC

SOLUTIONS

SAMPLE QUESTION PAPER

CBSE Class X Examination

10

Section 'A'

[1 × 20 = 20]

- | | | | |
|--------|---------|--------|--------|
| 1. (c) | 2. (b) | 3. (a) | 4. (c) |
| 5. (c) | 6. (d) | 7. (b) | 8. (c) |
| 9. (b) | 10. (d) | | |

Fill in the blanks

11. 8 cm


Explanation

$$AC^2 = (4\sqrt{2})^2 + (4\sqrt{2})^2$$

$$AC^2 = 32 + 32$$


$$AC^2 = 64$$

$$AC = 8 \text{ cm}$$



12. $x = \pm 3$

Explanation



Applying distance formula

$$OA = \sqrt{(x-0)^2 + (4-0)^2}$$

$$5 = \sqrt{x^2 + 16}$$

$$25 = x^2 + 16$$

$$x^2 = 9$$

$$x = \pm 3$$

13. $A = 45^\circ$

Explanation

$$\tan(A + B) = \sqrt{3} = \tan 60^\circ$$

$$\therefore A + B = 60^\circ$$

... (i)

$$\text{Similarly } \tan(A - B) = \frac{1}{\sqrt{3}} = \tan 30^\circ$$

$$\therefore A - B = 30^\circ$$


... (ii)

$$\text{Solving (i) and (ii) } A = 45^\circ$$

14. $A = 60^\circ$

Explanation

In $\triangle ABC$



$$\cos A = \frac{AB}{AC}$$

$$\cos A = \frac{3}{6} = \frac{1}{2} = \cos 60^\circ$$

$$\therefore A = 60^\circ$$

15. 1

Explanation

Given $\cos \alpha = \sin \alpha$

Divide by $\cos \alpha$

$$\Rightarrow \frac{\cos \alpha}{\cos \alpha} = \frac{\sin \alpha}{\cos \alpha}$$

$$\Rightarrow 1 = \tan \alpha$$

$$\therefore 5\alpha = 45$$

$$\text{Now } \tan 5\alpha = \tan 45^\circ = 1$$

Answer the following.

16. Let radius of circle bar

Circumference = Area of circle

$$2\pi r = \pi r^2$$

$$\Rightarrow r = 2$$

$$\therefore \text{Diameter} = 4$$

17. Volume of a ball with radius r

= Volume of 27 balls with radius r_1 .

$$\Rightarrow \frac{4}{3}\pi r^3 = 27 \times \frac{4}{3}\pi r_1^3$$

$$\Rightarrow \frac{r^3}{r_1^3} = \frac{27}{1}$$

$$\Rightarrow \left(\frac{r}{r_1}\right)^3 = \left(\frac{3}{1}\right)^3$$

$$\therefore r : r_1 = 3 : 1$$

$$18. \text{Sum of zeroes } S = \frac{3}{5} + \left(\frac{-1}{2}\right) = \frac{1}{10}$$

$$\text{Product of zeroes} = P = \frac{3}{5} \times \left(-\frac{1}{2}\right) = -\frac{3}{10}$$

\therefore Required polynomial is $x^2 - 8x + P = 0$

$$\Rightarrow x^2 - \frac{1}{10}x - \frac{3}{10} = 0$$

$$\text{or } 10x^2 - x - 3 = 0$$

19. Given equation is $9x^2 + 8kx + 16 = 0$

Here $a = 9$, $b = 8k$ and $c = 16$

For equal roots $b^2 - 4ac = 0$

$$(8k)^2 - 4 \times 9 \times 16 = 0$$

$$64k^2 - 64 \times 9 = 0$$

$$64(k^2 - 9) = 0$$

$$k^2 = 9$$

$$\text{or } k = \pm 3$$

$$20. S_n = 3x^2 + 5x$$

$$S_1 = 3 + 5 = 8 = a_1$$

$$S_2 = 3(2)^2 + 5(2) = 22 = a_1 + a_2$$

$$\therefore a_2 = 22 - 8 = 14$$

Now, the A.P. is 8, 14, ... 164

Here $a = 8$, $d = 6$ and $a_n = 164$

$$a_n = a + (n - 1)d$$

$$164 = 8 + (n - 1)6$$

$$\frac{156}{6} = n - 1$$

$$n = 27$$

Section 'B'

21. No, because $6^n = (2 \times 3)^n = 2^n \times 3^n$, so the only primes in the factorisation of 6^n are 2, 3 and not 5. [1]

Hence, it cannot end with the digit 5. [1]

$$22. ax + 3y = 5$$

$$a_1 = a, b_1 = 3, c_1 = 5$$

$$4x + 3ay = 10$$

$$a_2 = 4, b_2 = 3a, c_2 = 10$$

For infinitely many solutions $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

$$\frac{a}{4} = \frac{3}{3a} = \frac{5}{10} \quad [1]$$

$$\Rightarrow \frac{a}{4} = \frac{5}{10}$$

$$\Rightarrow a \times 10 = 5 \times 4$$

$$\Rightarrow a = \frac{20}{10}$$

$$\Rightarrow a = 2 \quad [1]$$

$$23. a + 3d = 0 \Rightarrow a = -3d$$

$$a_{25} = a + 24d = 21d \quad [1]$$

$$3a_{11} = 3(a + 10d)$$

$$= 3(7d) = 21d \quad [1]$$

[CBSE Marking Scheme, 2016]

CBSE Topper's Answer, 2016

you have,
 ~~$a_4 = 0$~~
 $a + 3d = 0$ $[a + (n-1)d = a_n]$
 $3d = -a$
 $a - 3d = a$ --- (1)

Now,
 ~~$a_{25} = a + 24d$~~ $[a + (n-1)d = a_n]$
 $-3d + 24d$
 $= 21d$ $\text{--- (2)} \quad (\text{Putting value of } 'a' \text{ from eq (1)})$
 $a_{11} = a + 10d$
 $-3d + 10d$
 $= 7d$ $\text{--- (2)} \quad (a = -3d)$
From eq (2) & eq (3)
 $\frac{a_{25}}{a_{11}} = \frac{3a_{11}}{3a_{11}}$
Derna Prasad.

24. Let the coordinates of points P and Q be $(0, b)$ and $(a, 0)$ resp.

$$\therefore \frac{a}{2} = 2 \Rightarrow a = 4 \quad [1]$$

$$\frac{b}{2} = -5 \Rightarrow b = -10 \quad [1]$$

$$\therefore P(0, -10) \text{ and } Q(4, 0) \quad [1]$$

[CBSE Marking Scheme, 2017]

25. (i) Probability of getting a doublet when two different dice are tossed. then

Total events $n(S) = 6 \times 6 = 36$

Favourable events i.e., to getting doublets

$$E = \{(1, 1), (2, 2), (3, 3), \\ (4, 4), (5, 5), (6, 6)\}$$

$$\therefore n(E) = 6$$

Probability of getting a doublet

$$= \frac{n(E)}{n(S)} = \frac{6}{36} = \frac{1}{6}$$

$$\therefore \text{Probability of getting a doublet} = \frac{1}{6} \quad [1]$$

- (ii) Favourable events of getting a sum 10, of the numbers on the two dice

$$E = \{(4, 6), (5, 5), (6, 4)\}$$

$$\therefore n(E) = 3$$

Probability of getting a sum 10, of the numbers on the two dice

$$= \frac{n(E)}{n(S)} = \frac{3}{36} = \frac{1}{12} \quad [1]$$

26. No. of all possible outcomes $= 6^2 = 36$

No. of favourable outcomes $= 26 \quad [1]$

$(4, 2) (4, 3) (4, 5) (5, 1) (5, 2) (5, 3) (3, 5)$
 $(6, 1) (6, 2) (1, 1) (1, 1) (1, 2) (1, 3) (1, 4)$
 $(1, 5) (1, 6) (2, 1) (2, 4) (2, 5) (2, 6) (3, 1)$
 $(3, 2) (3, 3) (3, 4) (2, 5) (4, 1)$

$\therefore P(\text{Product appears in less than } 18)$

$$= \frac{26}{36} = \frac{13}{18} \quad [1]$$

Section ‘C’

27. Let us assume, to contrary that $4 - 3\sqrt{2}$ is a rational number.

$$\therefore 4 - 3\sqrt{2} = \frac{p}{q}, q \neq 0 \text{ and } p, q \in \mathbb{Z} \quad [1]$$

$$\Rightarrow -3\sqrt{2} = \frac{p}{q} - 4$$

$$\Rightarrow -3\sqrt{2} = \frac{p - 4q}{q}$$

$$\Rightarrow \sqrt{2} = \frac{p - 4q}{-3q}$$

$$\Rightarrow \sqrt{2} = \frac{4q - p}{3q} = \frac{\text{Integer}}{\text{Integer}} \quad [1]$$

$$\Rightarrow \sqrt{2} = \text{a rational number}$$

But this contradicts the fact that $\sqrt{2}$ is irrational.

Hence, $4 - 3\sqrt{2}$ is an irrational number. [1]

28. Here $4\sqrt{3} \times (-2\sqrt{3}) = -24$, and $8 \times (-3) = -24$ and $8 + (-3) = 5$.

$$\therefore 4\sqrt{3}x^2 + 5x - 2\sqrt{3} = 0 \quad [1]$$

$$\Rightarrow 4\sqrt{3}x^2 + 8x - 3x - 2\sqrt{3} = 0$$

$$\Rightarrow 4x(\sqrt{3}x + 2) - \sqrt{3}(\sqrt{3}x + 2) = 0$$

$$\Rightarrow (4x - \sqrt{3})(\sqrt{3}x + 2) = 0$$


$$\Rightarrow \sqrt{3}x + 2 = 0 \text{ or } 4x - \sqrt{3} = 0$$

$$\Rightarrow x = \frac{-2}{\sqrt{3}} \text{ or } x = \frac{\sqrt{3}}{4} \quad [1]$$

$$\Rightarrow x = \left(\frac{-2}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} \right) = \frac{-2\sqrt{3}}{3} \text{ or } x = \frac{\sqrt{3}}{4}. \quad [1]$$

Hence, $\frac{-2\sqrt{3}}{3}$ and $\frac{\sqrt{3}}{4}$ are the roots of the given equation.

- 29.



ar ΔABC

$$\begin{aligned} &= \frac{1}{2} \left[x_1(y_2 - y_3) + x_2(y_3 - y_1) + x_3(y_1 - y_2) \right] \\ &= \frac{1}{2} [-5(-5+6) - 4(-6-7) - 1(7+5)] \\ &\quad = \frac{1}{2} |(-5+52-12)| \\ &\text{ar } \Delta ABC = \frac{1}{2} |35| = \frac{35}{2} \text{ unit}^2 \quad [1] \end{aligned}$$

ar ΔACD

$$\begin{aligned} &= \frac{1}{2} \left[x_1(y_3 - y_4) + x_3(y_4 - y_1) + x_4(y_1 - y_3) \right] \\ &\quad = \\ &\quad = \frac{1}{2} [-5(-6-5) + (-1)(5-7) + 4(7+6)] \end{aligned}$$

ar ΔACD

$$= \frac{1}{2} |(55+2+52)| = \frac{109}{2} \text{ unit}^2 \quad [1]$$

$$\text{ar of quadrilateral } ABCD = \text{ar } \Delta ABC + \text{ar } \Delta ACD$$

$$\begin{aligned} &= \frac{35}{2} + \frac{109}{2} \\ &= \frac{144}{2} = 72 \text{ unit}^2 \end{aligned}$$

$$\therefore \text{ar of quadrilateral } ABCD = 72 \text{ unit}^2 \quad [1]$$

OR

Let P (5, -2), Q (6, 4) and R (7, -2) be the given points.

\because Distance between (x_1, y_1) and (x_2, y_2)

$$= \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} \quad [\text{by distance formula}]$$

$$\begin{aligned} \therefore PQ &= \sqrt{(6-5)^2 + (4+2)^2} \\ &= \sqrt{(1)^2 + (6)^2} = \sqrt{1+36} = \sqrt{37} \text{ units} \quad [1] \end{aligned}$$


$$\begin{aligned} QR &= \sqrt{(7-6)^2 + (-2-4)^2} \\ &= \sqrt{(1)^2 + (-6)^2} = \sqrt{1+36} = \sqrt{37} \text{ units} \quad [1] \end{aligned}$$

$$\begin{aligned} \text{and PR} &= \sqrt{(7-5)^2 + (-2+2)^2} \\ &= \sqrt{2^2 + 0} = 2 \text{ units} \quad [1] \end{aligned}$$

$$\therefore PQ = QR \neq PR$$

$\therefore \Delta PQR$ is an isosceles triangle.

30.



[1]

As ΔACE and ΔDCP are equilateral triangles
 $\therefore \Delta ACE \sim \Delta DCP$ (AAA)

$$\begin{aligned} \frac{\text{ar } \Delta ACE}{\text{ar } \Delta DCP} &= \left(\frac{AC}{DC} \right)^2 \quad [1] \\ &= \left(\frac{\sqrt{2}a}{a} \right)^2 \\ &= \frac{2}{1} \end{aligned}$$

$$\therefore \Delta ACE = 2 \text{ ar } \Delta DCP \quad [1]$$

OR

Here, Since $PQ \parallel BC$ and PQ divides ΔABC into two equal parts, So $\Delta APQ \sim \Delta ABC$

$$\begin{aligned} &\therefore \frac{\text{ar } (\Delta APQ)}{\text{ar } (\Delta ABC)} = \frac{AP^2}{AB^2} \\ &\text{or, } \frac{1}{2} = \frac{AP^2}{AB^2} \quad [1] \\ &\text{or, } \frac{1}{\sqrt{2}} = \frac{AB - BP}{AB} \\ &\quad (\because AB = AP + BP) \\ &\text{or, } \frac{AB - BP}{AB} = \frac{1}{\sqrt{2}} \end{aligned}$$

$$\text{or, } 1 - \frac{BP}{AB} = \frac{1}{\sqrt{2}} \quad [1]$$

$$\text{or, } \frac{BP}{AB} = 1 - \frac{1}{\sqrt{2}} = \frac{\sqrt{2}-1}{\sqrt{2}}$$

$$\therefore BP : AB = (\sqrt{2}-1) : \sqrt{2} \quad [1]$$

31. Let the radius of circle be r cm. Then,

$$OA = OT = r \text{ cm.}$$

Since, PT is a tangent to circle at T and OT is a radius.

$$\text{So, } OT \perp PT$$

$$\therefore \angle OTP = 90^\circ \quad [1]$$

In right angled $\triangle OTP$,

$$OP^2 = OT^2 + PT^2 \quad \dots(i)$$

[by Pythagoras theorem]

$$\Rightarrow (PA + OA)^2 = OT^2 + 6^2$$

$$\Rightarrow (3+r)^2 = r^2 + 36$$

[from Eq. (i) and PA = 3 cm, PT = 6 cm, (given)]

$$\Rightarrow r^2 + 6r + 9 - r^2 - 36 = 0 \quad [1]$$

$$[\because (a+b)^2 = a^2 + b^2 + 2ab]$$

$$\Rightarrow 6r - 27 = 0 \Rightarrow r = \frac{27}{6} = 4.5$$

Hence, the radius of the circle is 4.5 cm [1]

$$32. \text{ To Prove } (\cot \theta - \operatorname{cosec} \theta)^2 = \frac{1-\cos \theta}{1+\cos \theta}$$

$$\begin{aligned} \text{LHS} &= (\cot \theta - \operatorname{cosec} \theta)^2 \\ &= \left(\frac{\cos \theta}{\sin \theta} - \frac{1}{\sin \theta} \right)^2 = \left(\frac{\cos \theta - 1}{\sin \theta} \right)^2 \quad [1] \\ &= \frac{(1-\cos \theta)^2}{\sin^2 \theta} \quad [1] \\ &\quad (\because \sin^2 \theta + \cos^2 \theta = 1) \end{aligned}$$

CBSE Topper's Answer, 2016

Q3. Here, Internal diameter of a hemispherical bowl = 36 cm
 $\Rightarrow \text{radius } R = 18 \text{ cm}$

AQ
The liquid of the bowl is filled into cylindrical bottles of diameter 6 cm i.e. of radius 3 cm
Also, 10% liquid is wasted.

Now, vol^m of the bowl = $\frac{2}{3} \times \pi \times 18^3 = \frac{2 \times \pi \times 18 \times 18 \times 18}{3}$
 $= (12 \times 324 \pi) \text{ cm}^3$

$$\begin{aligned} &= \frac{(1-\cos \theta)^2}{(1-\cos^2 \theta)} \\ &= \frac{(1-\cos \theta)(1-\cos \theta)}{(1-\cos \theta)(1+\cos \theta)} = \frac{1-\cos \theta}{1+\cos \theta} \quad [1] \\ &= \text{RHS} \quad \text{Hence Proved.} \end{aligned}$$

OR

We have:

$$\begin{aligned} \text{LHS} &= \frac{\sin \theta - 2 \sin^3 \theta}{2 \cos^3 \theta - \cos \theta} \\ &= \frac{\sin \theta(1 - 2 \sin^2 \theta)}{\cos \theta(2 \cos^2 \theta - 1)} \\ &= \tan \theta \cdot \frac{[1 - 2(1 - \cos^2 \theta)]}{(2 \cos^2 \theta - 1)} \\ &\quad [\because \sin^2 \theta = 1 - \cos^2 \theta] \\ &= \tan \theta \cdot \frac{(2 \cos^2 \theta - 1)}{(2 \cos^2 \theta - 1)} \\ &= \tan \theta = \text{RHS.} \end{aligned}$$

33. Volume of liquid in the bowl

$$= \frac{2}{3} \cdot \pi \cdot (18)^3 \text{ cm}^3$$

$$\text{Volume, after wastage} = \frac{2\pi}{3} \cdot (18)^3 \cdot \frac{90}{100} \text{ cm}^3$$

$$\text{Volume of liquid in 72 bottles} = \pi (3)^2 \cdot h \cdot 72 \text{ cm}^3$$

$$\Rightarrow h = \frac{\frac{2}{3} \pi (18)^3 \cdot \frac{9}{10}}{\pi (3)^2 \cdot 72} = 5.4 \text{ cm}$$

[3] [CBSE Marking Scheme, 2015]

~~liquid wasted = $\frac{10 \times 12 \times 3\pi}{100} = (1944\pi) \text{ cm}^3$~~

~~Note, liquid used to fill the bottles = $\frac{(3888\pi - 1944\pi)}{5} \text{ cm}^3$~~

~~= $\frac{(19440\pi - 1944\pi)}{5} \text{ cm}^3$~~

~~= $\frac{17496\pi}{5} \text{ cm}^3$~~

Now, height of each bottle = $\frac{19440\pi}{5 \times 3\pi \times 2 \times 3 \times 72} = \frac{27}{5} \text{ cm}$

= 5.4 cm

OR

Volume of water in the cylindrical tub
= volume of the tub

$$= \pi r^2 h = \left(\frac{22}{7} \times 5 \times 5 \times 9.8 \right) \text{ cm}^3$$


$$= 770 \text{ cm}^3.$$

Volume of the solid immersed in the tub
= volume of the hemisphere + volume of the cone

$$= \left(\frac{2}{3} \pi r^3 \right) + \left(\frac{1}{3} \pi r^2 h \right)$$

$$= \left[\left(\frac{2}{3} \times \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times \frac{7}{2} \right) + \left(\frac{1}{3} \times \frac{22}{7} \times \frac{7}{2} \times \frac{7}{2} \times 5 \right) \right] \text{ cm}^3$$

$$= \left(\frac{539}{6} + \frac{385}{6} \right) \text{ cm}^3 = \left(\frac{924}{6} \right) \text{ cm}^3 = 154 \text{ cm}^3.$$



Volume of water left in the tub

$$= (\text{volume of the tub}) - (\text{volume of solid immersed in the tub})$$

$$= (770 - 154) \text{ cm}^3 = 616 \text{ cm}^3.$$

34. Let assumed mean, $a = 35$ and given $h = 10$.

| Class | x_i (Class Marks) | $u_i = \frac{x_i - a}{h}$ | f_i | $f_i u_i$ |
|---------|------------------------|---------------------------|-----------------|----------------------|
| 0 – 10 | 5 | -3 | 5 | -15 |
| 10 – 20 | 15 | -2 | 13 | -26 |
| 20 – 30 | 25 | -1 | 20 | -20 |
| 30 – 40 | 35 | 0 | 15 | 0 |
| 40 – 50 | 45 | 1 | 7 | 7 |
| 50 – 60 | 55 | 2 | 5 | 10 |
| | Total | | $\sum f_i = 65$ | $\sum f_i u_i = -44$ |

$$\therefore \text{Mean}, \quad \bar{x} = a + \frac{\sum f_i u_i}{\sum f_i} \times h$$

$$= 35 + \frac{-44}{65} \times 10$$

$$= 35 - 6.76 = 28.24 \quad [1]$$

Section 'D'

35. Discriminant = $b^2 - 4ac = 36 - 4 \times 5 \times (-2)$
 $= 76 > 0$

So, the given equation has two distinct real roots

$$5x^2 - 6x - 2 = 0 \quad [1]$$

Multiplying both sides by 5.

$$(5x)^2 - 2 \times (5x) \times 3 = 10$$
 $\Rightarrow (5x)^2 - 2 \times (5x) \times 3 + 3^2 = 10 + 3^2$
 $\Rightarrow (5x - 3)^2 = 19 \quad [1]$
 $\Rightarrow 5x - 3 = \pm\sqrt{19}$

$$\Rightarrow x = \frac{3 \pm \sqrt{19}}{5} \quad [1]$$

Verification:

$$5\left(\frac{3+\sqrt{19}}{5}\right)^2 - 6\left(\frac{3+\sqrt{19}}{5}\right) - 2 \quad [1]$$

$$= \frac{9+6\sqrt{19}+19}{5} - \frac{18+6\sqrt{19}}{5} - \frac{10}{5} = 0$$

$$\text{Similarly, } 5\left(\frac{3-\sqrt{19}}{5}\right)^2 - 6\left(\frac{3-\sqrt{19}}{5}\right) - 2 = 0 \quad [1]$$

[CBSE Marking Scheme, 2018]

OR

Let the length of shorter side be x m.

∴ length of diagonal = $(x + 16)$ m [1]

and, length of longer side = $(x + 14)$ m [1]


∴ $x^2 + (x + 14)^2 = (x + 16)^2$ [1]

⇒ $x^2 - 4x - 60 = 0 \Rightarrow x = 10$ m

∴ length of sides are 10m and 24m. [1]

CBSE Topper's Solution, 2015

Let ABCD be a rectangle.
 Let the shorter side, BC = x cm
 Then, AC = $(x+16)$ cm
 And, AB = $(x+14)$ cm



Now, in right $\triangle ABC$, By Pythagoras theorem

$$AC^2 = AB^2 + BC^2$$

$$\text{Or, } (x+16)^2 = (x+14)^2 + x^2$$

$$\text{Or, } x^2 + 256 + 32x = x^2 + 196 + 28x + x^2$$

$$\text{Or, } 2x^2 + 28x + 196 - x^2 - 32x - 256 = 0$$

$$\text{Or, } x^2 - 4x - 60 = 0, \text{ which is a quad. eqn.}$$

$$\text{Or, } x^2 - 10x + 6x - 60 = 0$$

$$\text{Or, } x(x-10) + 6(x-10) = 0$$

$$\text{Or, } (x-10)(x+6) = 0$$

$$\text{Or, } x-10 = 0 \quad \text{Or, } x+6 = 0$$


$$\text{Or, } x = 10 \quad \text{Or, } x = -6$$

(3 invalid)

Now, BC = $x = 10$ cm

$$AB = x+14 = (10+14) \text{ m} = 24 \text{ m}$$

36. Let $AD = 3x$ cm and $DB = 2x$ cm.



Then,

$$AB = (AD + DB) = (3x + 2x) \text{ cm} = 5x \text{ cm.}$$

[1]

In $\triangle ABC$ and $\triangle DBE$, we have:

$$\angle CAB = \angle EDB \text{ (corresponding } \angle\text{s)}$$

$$\angle ACB = \angle DEB \text{ (corresponding } \angle\text{s)}$$

$\therefore \triangle ABC \sim \triangle DBE$ [by AA-similarity] [1]

$$\Rightarrow \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DBE)} = \frac{(AB)^2}{(DB)^2} = \frac{(5x)^2}{(2x)^2} = \frac{25}{4}$$

[1]

$$\Rightarrow \text{ar}(\triangle ABC) : \text{ar}(\triangle DBE) = 25 : 4.$$

OR


Given : $\triangle ABC \sim \triangle PQR$

To Prove :

$$\begin{aligned} \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} &= \left(\frac{AB}{PQ} \right)^2 \\ &= \left(\frac{BC}{QR} \right)^2 = \left(\frac{AC}{PR} \right)^2 \end{aligned}$$

Construction : Draw $AD \perp BC$ and $PE \perp QR$

Proof : $\triangle ABC \sim \triangle PQR$



$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR}$$

(Corresponding sides of similar triangles)
...(i)

In $\triangle ADB$ and $\triangle PEQ$,

$$\angle B = \angle Q \text{ (Proved)}$$

$$\angle ADB = \angle PEQ \quad [\text{each } 90^\circ]$$

$\therefore \triangle ADB \sim \triangle PEQ$ (AA similarity)

$$\text{or,} \quad \frac{AD}{PE} = \frac{AB}{PQ}$$

(Corresponding sides of similar triangles) ... (ii)

From eq. (i) and eq. (ii),

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AC}{PR} = \frac{AD}{PE}$$

...(iii)

$$\begin{aligned} \text{Now} \quad \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} &= \frac{\frac{1}{2} \times BC \times AD}{\frac{1}{2} \times QR \times PE} \\ &= \left(\frac{BC}{QR} \right) \times \left(\frac{AD}{PE} \right) \\ &= \frac{BC}{QR} \times \frac{BC}{QR} \end{aligned}$$

$$\text{or,} \quad \frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \frac{BC^2}{QR^2} \quad \dots (iv)$$


[from eq. (iii)]

From eq (iii) and eq (iv),

$$\frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle PQR)} = \left(\frac{AB}{PQ} \right)^2$$

$$= \left(\frac{BC}{QR} \right)^2 = \left(\frac{AC}{PR} \right)^2$$


37. CBSE Topper's Answer, 2015

Steps:

1. we draw $\triangle ABC$ with the given dimensions.
2. we draw a perpendicular to AC at D from B.
3. we draw a circle passing through B, C & D.
4. we draw tangent to the circle from A at B & E.
 $\therefore AB \text{ & } AE$ are required tangents.

[4]

38.



[1]

$$\frac{100}{x} = \tan 45^\circ = 1$$

$$\Rightarrow x = 100 \quad \dots(i) [1]$$


$$\frac{100}{y} = \tan 30^\circ = \frac{1}{\sqrt{3}}$$

$$\Rightarrow y = 100\sqrt{3} \quad \dots(ii) [1]$$


Distance between the cars = $x + y$

$$= 100(\sqrt{3} + 1) = 273.2 \text{ m} \quad [1]$$

CBSE Topper's Answer, 2017



39. Total Volume of cylinder = $\frac{22}{7} \times \frac{22}{10} \times \frac{42}{10} \times 10 \text{ cm}^3 = 554.40 \text{ cm}^3$



[1]

Volume of metal scooped out = $\frac{4}{3} \times \frac{22}{7} \times \left(\frac{42}{10}\right)^3 = 310.46 \text{ cm}^3$

[1]

\therefore Volume of rest of cylinder = $554.40 - 310.46 = 243.94 \text{ cm}^3$

[1]

If l is the length of wire, then


$$\frac{22}{7} \times \frac{7}{10} \times \frac{7}{10} \times l = \frac{24394}{100}$$

$$\Rightarrow l = 158.4 \text{ cm}$$

[1]

[CBSE Marking Scheme, 2017]

CBSE Topper's Answer, 2015



40. (i) By Formula Method :

| Classes | f | c.f. | |
|---------|-----|------|----------------|
| 0–20 | 6 | 6 | |
| 20–40 | 8 | 14 | |
| 40–60 | 10 | 24 | |
| 60–80 | 12 | 36 | ⇒ Median Class |
| 80–100 | 6 | 42 | |
| 100–120 | 5 | 47 | |
| 120–140 | 3 | 50 | |

[1]

$$\text{Median} = \frac{N}{2} \text{ th term} = \frac{50}{2} = 25 \text{th term}$$

Median class = 60 – 80

$$\text{Median} = l + \left(\frac{\frac{N}{2} - c.f.}{f} \right) \times h = 60 + \left(\frac{25 - 24}{12} \right) \times 20$$

$$= 60 + \frac{1}{12} \times 20 = 60 + \frac{5}{3} = \frac{185}{3} = 61.67$$


[½]

(ii)

| Classes | c.f. |
|---------------|------|
| More than 0 | 50 |
| More than 20 | 44 |
| More than 40 | 36 |
| More than 60 | 26 |
| More than 80 | 14 |
| More than 100 | 8 |
| More than 120 | 3 |

[1]

To draw ogive we take the indeces : (0, 50), (20, 44), (40, 36), (60, 26), (80, 14), (100, 8), (120, 3)



[1]

From graph,

$$\frac{N}{2} = \frac{50}{2} = 25$$

$$\text{Median} = 61.6$$

[½]

OR

Since the mode of the given series is 36 and maximum frequency 16 lies in the class 30–40, so the modal class is 30–40.

Let the missing frequency be x . Then

$$\therefore x_k = 30, f_k = 16, f_{k-1} = x, f_{k+1} = 12 \text{ and } h = 10.$$

Also, $M_o = 36$.

Using the formula, $M_o = x_k + \left\{ h \times \frac{(f_k - f_{k-1})}{(2f_k - f_{k-1} - f_{k+1})} \right\}$, we get:

$$30 + \left\{ 10 \times \frac{(16 - x)}{(32 - x - 12)} \right\} = 36$$

$$\Rightarrow \frac{10 \times (16 - x)}{(20 - x)} = 6 \Rightarrow 160 - 10x = 120 - 6x$$

$$\Rightarrow 4x = 40 \Rightarrow x = 10.$$

Hence, the missing frequency is 10.

MATHEMATICS – BASIC

SOLUTIONS

SAMPLE QUESTION PAPER

CBSE Class X Examination

11

Section ‘A’

[1 × 20 = 20]

- | | | | |
|--------|---------|--------|--------|
| 1. (a) | 2. (b) | 3. (c) | 4. (c) |
| 5. (a) | 6. (a) | 7. (b) | 8. (a) |
| 9. (d) | 10. (c) | | |

Fill in the blanks

11. $p = 4$

Explanation

$(2p + 1), 13, (5p - 3)$ are in A.P.

$$\begin{aligned} \text{Common difference} &= (5p - 3) - 13 \\ &= 13 - (2p + 1) \\ \Rightarrow 5p - 16 &= 12 - 2p \\ 7p &= 28 \\ p &= 4 \end{aligned}$$

12. 2139

Explanation

Here $a = 5$ and $d = 13 - 5 = 8$

$$\begin{aligned} a_x &= a + (n - 1)d \\ |8| &= 5 + (n - 1)8 \\ 176 &= 8(x - 1) \\ x - 1 &= 22 \\ x &= 23 \\ \therefore S_n &= \frac{n}{2}[a + a_n] \\ &= \frac{23}{2}[5 + 181] \\ &= \frac{23}{2} \times 186 = 2139 \end{aligned}$$

13. $x = \frac{\sqrt{2}}{\sqrt{3}}, \frac{\sqrt{2}}{\sqrt{3}}$

Explanation

$$\begin{aligned} 3x^2 - 2\sqrt{6}x + 2 &= 0 \\ 3x^2 - \sqrt{6}x - \sqrt{6}x + 2 &= 0 \\ 3x^2 - \sqrt{3}\sqrt{2}x - \sqrt{3}\sqrt{2}x + 2 &= 0 \\ \sqrt{3}x[\sqrt{3}x - \sqrt{2}] - \sqrt{2}[\sqrt{3}x - \sqrt{2}] &= 0 \\ (\sqrt{3}x - \sqrt{2})(\sqrt{3}x - \sqrt{2}) &= 0 \end{aligned}$$

$$x = \frac{\sqrt{2}}{\sqrt{3}}, \frac{\sqrt{2}}{\sqrt{3}}$$

14. Imaginary

Explanation

$$\begin{aligned} x^2 + x + 1 &= 0 \\ \text{Here } a &= 1, b = 1, c = 1 \\ D &= b^2 - 4ac \\ D &= 1 - 4 = -3 \\ D < 0 & \end{aligned}$$

15. LCM = 338

Explanation

$$\begin{aligned} \text{Given } N_1 &= 26 \\ N_2 &= 169 \end{aligned}$$

and H.C.F. = 13

$$\begin{aligned} \text{We know that } N_1 \times N_2 &= \text{HCF} \times \text{LCM} \\ \Rightarrow 26 \times 169 &= 13 \times \text{LCM} \\ \therefore \text{LCM} &= 338 \end{aligned}$$

Answer the following.

16. Let outer radius of path be R and inner radius be r .

According to question:

$$\frac{2\pi R}{2\pi r} = \frac{23}{22}$$

$$\Rightarrow \frac{R}{r} = \frac{23}{22} \quad \dots \text{(i)}$$

Also $R - r = 5\text{m}$

$$\Rightarrow R = (5 + r)\text{m}$$

Solving (i) and (ii)

$$\frac{5+r}{r} = \frac{23}{22}$$

$$110 + 22r = 23r$$

$$\Rightarrow r = 110 \text{ m}$$

\therefore Diameter = 220 m

17. Given $Q = 45^\circ$

$$r = 7 \text{ cm}$$

$$\text{Arc length } l = \frac{Q}{360} \times 2\pi r$$

$$l = \frac{45}{360} \times 2 \times \frac{22}{7} \times 7 \\ = 5.5 \text{ cm}$$

18. Curved surface area = $2\pi rh = 264$

$$\text{Volume} = \pi r^2 h = 264$$

$$\text{Dividing (i) } \div \text{ ii } \frac{2\pi rh}{\pi r^2 h} = \frac{264}{924}$$

$$\frac{2}{r} = \frac{2}{7}$$


$$\Rightarrow r = 7 \Rightarrow \text{Diameter} = 14$$

$$\text{from (i) } 2 \times \frac{22}{7} \times 7 \times h = 264$$

$$h = 6$$

$$\therefore \frac{\text{Diameter}}{\text{height}} = \frac{14}{6} = \frac{7}{3}$$

- 19.



In $\triangle ABC$

$$\tan 30^\circ = \frac{BC}{9}$$

$$\frac{1}{\sqrt{3}} = \frac{BC}{9} \Rightarrow BC = \frac{9}{\sqrt{3}}$$

$$BC = \frac{9}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = 3\sqrt{3} \text{ cm}$$

20. $\sin A = \cos B$

$$\sin A = \sin (90 - B)$$

$$\Rightarrow A = 90 - B$$

$$\Rightarrow A + B = 90^\circ$$

Section 'B'

21. $kx + y = k^2$ and $x + ky = 1$

$$\frac{a_1}{a_2} = \frac{k}{1}, \frac{b_1}{b_2} = \frac{1}{k}, \frac{c_1}{c_2} = \frac{k^2}{1} \quad [1]$$

For infinitely many solutions

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$$

$$\frac{k}{1} = \frac{1}{k} = \frac{k^2}{1} = k^2 = 1$$

$$k = \pm 1 \quad [1]$$

22. Let us assume to the contrary, that $3\sqrt{7}$ is an rational.

Then, there exist co-prime positive integers p and q such that

$$3\sqrt{7} = \frac{p}{q} \quad [1]$$

$\left[\because p, 3 \text{ and } q \text{ are integers} \right]$

$\left[\because \frac{p}{3q} \text{ is a rational number} \right]$

$$\Rightarrow \sqrt{7} = \frac{p}{3q}$$

$\Rightarrow \sqrt{7}$ is rational [1]

But this contradicts the fact that $\sqrt{7}$ is irrational.

This contradiction has arisen because of our incorrect assumption that $3\sqrt{7}$ is rational. So, we conclude that $3\sqrt{7}$ is an irrational number.

23. Here, first term (a) = 6

Common difference (d) of the given

$$\text{A.P.} = 13 - 6 = 7 \quad [1]$$

Let the given A.P. contains n terms, then

$$t_n = 216 \text{ (given)}$$

$$\Rightarrow a + (n-1)d = 216$$

$$\Rightarrow 6 + (n-1)7 = 216$$

$$\Rightarrow (n-1)7 = 210$$

$$\Rightarrow n-1 = 30$$

$$\Rightarrow n = 30 + 1 = 31$$

Thus, the given A.P. contains 31 terms. [1]

24. Let the point be A(3, 0), B(6, 4), C(-1, 3)
- $$AB = \sqrt{9+16} = 5,$$
- $$BC = \sqrt{49+1} = 5\sqrt{2},$$
- $$AC = \sqrt{16+9} = 5$$

$AB = AC$
and $AB^2 + AC^2 = BC^2$
 ΔABC isosceles right angled triangle.
[1½ + ½] [CBSE Marking Scheme, 2016]

CBSE Topper's Answer, 2016

Let $A = (3, 0)$; $B = (6, 4)$ and $C = (-1, 3)$.
 Applying distance formula –
 $AB = \sqrt{(6-3)^2 + (4-0)^2} = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$ unit
 $BC = \sqrt{(6+1)^2 + (4-3)^2} = \sqrt{7^2 + 1^2} = \sqrt{49+1} = \sqrt{50} = 5\sqrt{2}$ unit
 $AC = \sqrt{(3+1)^2 + (0-3)^2} = \sqrt{25} = 5$ unit.
 Since,
 $AB = AC = 5$ unit
 ΔABC is isosceles triangle.
 Also, $AB^2 + AC^2 = BC^2$
 $\Rightarrow 25 + 25 = 50 = BC^2 = (5\sqrt{2})^2 \Rightarrow AB^2 + AC^2 = BC^2$
 Hence, by converse of Pythagoras Theorem,
 $\Rightarrow \Delta ABC$ is right angled triangle.

25. (i) $P(\text{square number}) = \frac{8}{113}$ [1]
 (ii) $P(\text{multiple of 7}) = \frac{16}{113}$ [1]
- [CBSE Marking Scheme, 2018]

26. No. of possible outcomes = 100
 Perfect squares 4, 9, 16, 25, 36, 49, 64, 81, 100
 No. of favourable outcomes = 9
- (i) $P(\text{perfect square}) = \frac{9}{100}$ [1]
 (ii) $P(\text{odd numbers not less than 70}) = \frac{16}{100} = \frac{4}{25}$ [1]

$$\begin{aligned} \text{Also, } LCM + HCF &= 600 & [1] \\ \Rightarrow 14 \times HCF + HCF &= 600 \\ \Rightarrow 15 HCF &= 600 \\ \Rightarrow HCF &= 40 & [1] \\ \therefore LCM &= 14 \times 40 = 560 \end{aligned}$$

$$\begin{aligned} \text{Now, one number is 280} \\ \therefore 280 \times \text{Other number} &= 40 \times 560 \\ \Rightarrow \text{Other number} &= \frac{40 \times 560}{280} \\ &= 80 & [1] \end{aligned}$$

Section 'C'

27. According to the statement of the question, we have
 $LCM \text{ of two numbers} = 14 \times HCF \text{ of two numbers}$

28. Let the ten's and the units digit be y and x respectively.
 So, the number is $10y + x$.
 The number when digits are reversed is $10x + y$.
 $Now, 7(10y + x) = 4(10x + y) \Rightarrow 2y = x$... (i)
 $Also x - y = 3$... (ii)
 Solving (1) and (2), we get $y = 3$ and $x = 6$.
- [3] [CBSE Marking Scheme, 2018]


29. Let the given points be $A(-5, 2)$ and $B(9, -2)$ and let the required point be $P(0, y)$. Then,

$$\begin{aligned} PA = PB &\Rightarrow PA^2 = PB^2 \quad [1] \\ \Rightarrow (0+5)^2 + (y-2)^2 &= (0-9)^2 + (y+2)^2 \quad [1] \\ \Rightarrow 5^2 + (y-2)^2 &= (-9)^2 + (y+2)^2 \\ \Rightarrow 25 + (y-2)^2 &= 81 + (y+2)^2 \\ \Rightarrow (y+2)^2 - (y-2)^2 &= (25-81) \\ \Rightarrow 8y &= -56 \Rightarrow y = -7. \end{aligned}$$

Hence, the required point is P(0, -7).

OR

Let PA : AQ = k : 1



$$\therefore \frac{2+3k}{k+1} = \frac{24}{11} \quad [1]$$

$$\Rightarrow k = \frac{2}{9}$$

Hence the ratio is 2 : 9. [1]


$$\text{Therefore } y = \frac{-18+14}{11} = \frac{-4}{11} \quad [1]$$

[CBSE Marking Scheme, 2017]

CBSE Topper's Answer 2017

$$\begin{aligned} &\text{Given } P(2, -2), Q(3, 7) \quad \text{Here } x_1 = 2, y_1 = -2 \\ &\text{Using section formula, } (\frac{24}{11}, y) = \left(\frac{3m+2n}{m+n}, \frac{7m-2n}{m+n} \right) \quad \text{--- (1)} \\ &\Rightarrow \frac{24}{11} = \frac{3m+2n}{m+n} \\ &24m+24n = 33m+22n \\ &2n = 9m \\ &\frac{2}{9} = \frac{m}{n} \\ &\therefore \text{The given point divides the line segment in ratio } 2:9. \\ &\text{Taking } m=2 \text{ and } n=9, \\ &y = \frac{7m-2n}{m+n} \quad (\text{from (1)}) \\ &y = \frac{7(2)-2(9)}{2+9} \\ &y = \frac{14-18}{11} \\ &y = -\frac{4}{11} \end{aligned}$$

30. $\angle SQN = \angle TRM$ (CPCT as $\triangle NSQ \cong \triangle MTR$)




Since, $\angle P + \angle 1 + \angle 2 = \angle P + \angle PQR + \angle PRQ$
(Angle sum property)

$$\Rightarrow \angle 1 + \angle 2 = \angle PQR + \angle PRQ$$

$$\begin{aligned} &\Rightarrow 2\angle 1 = 2\angle PQR \\ &\text{(as } \angle 1 = \angle 2 \text{ and } \angle PQR = \angle PRQ) \\ &\angle 1 = \angle PQR \\ &\angle 2 = \angle PRQ \\ &\text{And } \angle SPT = \angle QPR \text{ (common)} \\ &\triangle PTS \sim \triangle PRQ \\ &\text{(By AAA similarity criterion)} \\ &[\text{CBSE Marking Scheme, 2015}] [1+1+1] \end{aligned}$$

OR

Let AB be the street and let C be the foot of the ladder. Let D and E be the given windows such that AD = 9 m and BE = 12 m.



Then, CD and CE are the two positions of the ladder.

Clearly, $\angle CAD = 90^\circ$, $\angle CBE = 90^\circ$ and $CD = CE = 15 \text{ m}$.

From right $\triangle CAD$, we have:

$$CD^2 = AC^2 + AD^2$$

[by Pythagoras' theorem]

$$\Rightarrow AC^2 = (CD^2 - AD^2) \\ = [(15)^2 - (9)^2] \text{ m}^2 \\ = (225 - 81) \text{ m}^2$$

$$\Rightarrow AC = \sqrt{144} \text{ m} = 12 \text{ m}$$

from right $\triangle CBE$, we have

$$CE = CB^2 + BE^2 \text{ (by Pythagoras' theorem)} \\ \Rightarrow CB^2 = (CE^2 - BE^2) = [(15)^2 - (12)^2] \text{ m}^2 \\ = (225 - 144) \text{ m}^2 = 81 \text{ m}^2$$

$$\Rightarrow CB = \sqrt{81} \text{ m} = 9 \text{ m.}$$

Width of the street = $(AC + CB)$

$$= (12 + 9) \text{ m} = 21 \text{ m.}$$

31. LHS = $\frac{\sin \theta - \cos \theta}{\sin \theta + \cos \theta} + \frac{\sin \theta + \cos \theta}{\sin \theta - \cos \theta} = \frac{(\sin \theta - \cos \theta)^2 + (\sin \theta + \cos \theta)^2}{\sin^2 \theta - \cos^2 \theta}$

$$= \frac{(\sin^2 \theta + \cos^2 \theta) - 2 \sin \theta \cos \theta + (\sin^2 \theta + \cos^2 \theta) + 2 \sin \theta \cos \theta}{\sin^2 \theta - (1 - \sin^2 \theta)}$$

$$= \frac{1 + 1}{\sin^2 \theta - 1 + \sin^2 \theta} = \frac{2}{2 \sin^2 \theta - 1} = \text{RHS}$$

[1 + 1 + 1][CBSE Marking Scheme, 2015]

OR

To prove: $\frac{1}{\operatorname{cosec} A - \cot A} - \frac{1}{\sin A} = \frac{1}{\sin A} - \frac{1}{\operatorname{cosec} A + \cot A}$

$$\frac{1}{\operatorname{cosec} A - \cot A} - \frac{1}{\sin A} = \frac{1}{\sin A} - \frac{1}{\operatorname{cosec} A + \cot A}$$

[1]

or $\frac{1}{\operatorname{cosec} A - \cot A} + \frac{1}{\operatorname{cosec} A + \cot A} = \frac{1}{\sin A} + \frac{1}{\sin A}$

or $\frac{1}{\operatorname{cosec} A - \cot A} + \frac{1}{\operatorname{cosec} A + \cot A} = \frac{2}{\sin A}$

[1]

L.H.S. = $\frac{1}{\operatorname{cosec} A - \cot A} + \frac{1}{\operatorname{cosec} A + \cot A}$ or $\frac{\operatorname{cosec} A + \cot A + \operatorname{cosec} A - \cot A}{(\operatorname{cosec} A - \cot A)(\operatorname{cosec} A + \cot A)}$

or $\frac{2\operatorname{cosec} A}{\operatorname{cosec}^2 A - \cot^2 A}$ or $\frac{\frac{2}{\sin A}}{1} = \frac{2}{\sin A} = \text{R.H.S.}$

[1]

Hence Proved

32. Let r cm and R m be the radii of inner and outer boundaries.

Then, $2\pi r = 437$ and $2\pi R = 503$

$$\Rightarrow r = \frac{437}{2\pi} \text{ and } R = \frac{503}{2\pi}.$$

[1]

Width of the track = $(R - r)$ m


$$= \left(\frac{503}{2\pi} - \frac{437}{2\pi} \right) \text{m} = \frac{1}{2\pi} \times (503 - 437) \text{ m}$$

[1]

$$= \left(\frac{1}{2} \times \frac{7}{22} \times 66 \right) \text{m}$$

$$= 10.5 \text{ m.}$$

[1]



33. Largest possible diameter of hemisphere = 10 cm

\therefore radius = 5 cm

$$\text{Total surface area} = 6(10)^2 + 3.14 \times (5)^2$$

$$\text{Cost of painting} = \frac{678.5 \times 5}{100} = \frac{\text{₹ } 3392.50}{100} = \text{₹ } 33.9250 = \text{₹ } 33.93$$

[3] [CBSE Marking Scheme, 2015]


CBSE Topper's Answer, 2015

Here, side of a cubical block = $a = 10 \text{ cm}$

A hemisphere surrounds the cube.

\therefore Diameter (largest) of hemisphere = 10 cm

$$\Rightarrow \text{radius } (r) = 5 \text{ cm}$$



NOW, T.S.A of solid = (T.S.A of cube - area of base of hemisphere) + C.S.A of hemisphere

$$= 6a^2 - \pi r^2 + 2\pi r^2$$

$$= 6a^2 + \pi r^2 = 6(10)^2 + \pi(5)^2$$

$$= 600 + 25 \times 3.14$$

$$= 600 + 25 \times 3.14 \times 157$$

$$= 1200 + 157 = 1357 \text{ cm}^2$$

$$= \frac{1357}{2} = 678.5 \text{ cm}^2$$

NOW, Rate of painting = ₹ 5/100 cm²

$$\therefore \text{Cost} \rightarrow = \text{₹ } \frac{5 \times 678.5}{100}$$

$$= \text{₹ } \frac{3392.50}{20}$$

$$= \text{₹ } 169.625$$

$$= \text{₹ } \frac{169.625}{20} = \text{₹ } 8.48125$$

$$= \text{₹ } \frac{8.48125}{20} = \text{₹ } 0.4240625$$

$$= \text{₹ } 0.4240625 \times 100 = \text{₹ } 42.40625$$

$$= \text{₹ } 42.40625 \times 20 = \text{₹ } 848.125$$

$$= \text{₹ } 848.125 \times 20 = \text{₹ } 16962.50$$

$$= \text{₹ } 16962.50 \times 20 = \text{₹ } 339250$$

$$= \text{₹ } 339250 \times 20 = \text{₹ } 678500$$

$$= \text{₹ } 678500 \times 20 = \text{₹ } 1357000$$

$$= \text{₹ } 1357000 \times 20 = \text{₹ } 2714000$$

$$= \text{₹ } 2714000 \times 20 = \text{₹ } 5428000$$

$$= \text{₹ } 5428000 \times 20 = \text{₹ } 10856000$$

OR

Volume of cuboid = $4.4 \times 2.6 \times 1 \text{ m}^3$

Inner and outer radii of cylindrical pipe = 30 cm, 35 cm

$$\therefore \text{Volume of material used} = \frac{\pi}{100^2} (35^2 - 30^2) \times h \text{ m}^3 = \frac{\pi}{100^2} \times 65 \times 5 h$$

$$\text{Now } \frac{\pi}{100^2} \times 65 \times 5 h = 4.4 \times 2.6$$

$$\Rightarrow h = \frac{7 \times 4.4 \times 2.6 \times 100 \times 100}{22 \times 65 \times 5}$$

$$\Rightarrow h = 112 \text{ m}$$

[3] [CBSE Marking Scheme, 2017]

CBSE Topper's Answer, 2017

For the hollow cylindrical pipe,

$$r = 30 \text{ cm} \quad \text{and} \quad R = 30 + 5 = 35 \text{ cm}$$

Let its length be h .

volume of the 2 is same.

$$\therefore 4.4 \times 100 \times 2.6 \times 100 \times 100 = \pi h (R^2 - r^2)$$

$$440 \times 260 \times 100 = 22 \times h \times (35^2 - 30^2)$$

$$440 \times 260 \times 100 = 22 \times h \times 65 \times 5$$

$$\frac{7 \times 440}{22} \times \frac{260}{65} \times \frac{100}{5} = h$$

$$7 \times 20 \times 4 \times 20 = h$$

\therefore pipe is 11200 cm or 112 m long

34.

| C.I. | x | f | $u_i = \frac{x-f}{h}$ | $f_i u_i$ |
|-------|----|--------|-----------------------|-----------|
| 0–10 | 5 | 5 | -3 | -15 |
| 10–20 | 15 | x | -2 | -2x |
| 20–30 | 25 | 10 | -1 | -10 |
| 30–40 | 35 | 12 | 0 | 0 |
| 40–50 | 45 | 7 | 1 | 7 |
| 50–60 | 55 | 8 | 2 | 16 |
| Total | | $42+x$ | | $-2x-2$ |

$$\begin{aligned}
 A &= \text{Assumed mean} \\
 &= \text{mid point of class } 30 - 40 \\
 &= 35 \\
 \text{Mean} &= A + \frac{\sum f_i u_i}{\sum f_i} \times h \\
 \Rightarrow 31.4 &= 35 + \frac{-2x - 2}{42 + x} \times 10 \\
 \Rightarrow (2x + 2)10 &= (42 + x)(3.6) \\
 \Rightarrow 20x + 20 &= 151.2 + 3.6x \\
 16.4x &= 131.2 \\
 x &= 8 \quad [1]
 \end{aligned}$$

$$\begin{aligned}
 \text{B's one day work} &= \frac{1}{x} \quad [1] \\
 \text{A's one day work} &= \frac{1}{x-6} \quad [1] \\
 \text{and (A + B)'s one day work} &= \frac{1}{4} \\
 \text{According to the question, } \frac{1}{x} + \frac{1}{x-6} &= \frac{1}{4} \\
 \text{or } x^2 - 14x + 24 &= 0 \quad [1] \\
 \text{or } x^2 - 12x - 2x + 24 &= 0 \\
 \text{or } x(x-12) - 2(x-12) &= 0 \\
 \text{or } (x-12)(x-2) &= 0 \\
 \text{or } x = 12 \text{ or } x = 2 \\
 \text{But } x \text{ cannot be less than 6. So } x = 12 \quad [1] \\
 \text{Hence, B can finish the work in 12 days.}
 \end{aligned}$$

Section 'D'

35. Suppose B alone finish the work in x days and A alone takes $(x-6)$ days.

CBSE Topper's Answer, 2017

Let B complete a work in x days.
 Then A takes $x-6$ days to complete it.
 Together they complete it in 4 days.
 According to work done per day,

$$\begin{aligned}
 \frac{1}{x-6} + \frac{1}{x} &= \frac{1}{4} \\
 \frac{x + x-6}{x(x-6)} &= \frac{1}{4} \\
 4(x-6) &= x(x-6) \\
 8x - 24 &= x^2 - 6x \\
 x^2 - 14x + 24 &= 0 \\
 x(x-12) - 2(x-12) &= 0 \\
 (x-2)(x-12) &= 0 \\
 \therefore x = 2 \text{ or } 12. \\
 x = 2 \text{ is not possible because then } x-6 &\text{ is } \in 4 \\
 \therefore x = 12. \\
 \text{So, B takes 12 days to finish the work.}
 \end{aligned}$$


OR

The given equation is $2x^2 + kx + 3 = 0$.
 This is of the form $ax^2 + bx + c = 0$, where
 $a = 2, b = k$ and $c = 3$.

$$\begin{aligned}
 \therefore D &= (b^2 - 4ac) = (k^2 - 4 \times 3) \times 2 \\
 &= (k^2 - 24).
 \end{aligned}$$

For real and equal roots, we must have:

$$\begin{aligned}
 D = 0 &\Rightarrow k^2 - 24 = 0 \Rightarrow k^2 = 24 \\
 \Rightarrow k &= \pm\sqrt{24} = \pm 2\sqrt{6}. \\
 \text{Hence, } 2\sqrt{6} \text{ and } -2\sqrt{6} &\text{ are the required values of } k.
 \end{aligned}$$



Given : $AB \perp BC$

Construction : Draw $BE \perp AC$

To Prove : $AB^2 + BC^2 = AC^2$

Proof : In $\triangle AEB$ and $\triangle ABC$ $\angle A = \angle A$
(Common)

$\angle E = \angle B$ (each 90°)

$\triangle AEB \sim \triangle ABC$
(By AA similarity)

or

$$\frac{AE}{AB} = \frac{AB}{AC}$$

or

$$AB^2 = AE \times AC \quad \dots(i)$$

Now, in $\triangle CEB$ and $\triangle CBA$, $\angle C = \angle C$
(Common)

$\angle E = \angle B$ (each 90°)

$\triangle CEB \sim \triangle CBA$
(By AA similarity)

or

$$\frac{CE}{BC} = \frac{BC}{AC}$$

or

$$BC^2 = CE \times AC \quad \dots(ii) [1]$$

On adding eqns. (i) and (ii),


$$AB^2 + BC^2 = AE \times AC + CE \times AC$$

$$\text{or } AB^2 + BC^2 = AC(AE + CE)$$

$$\text{or } AB^2 + BC^2 = AC \times AC$$

$$\therefore AB^2 + BC^2 = AC^2 \quad [1]$$

(ii)



Given: ABCD is a rhombus,

Construction: Draw diagonals AC and BD

$$\therefore AO = OC = \frac{1}{2} AC$$

and $BO = OD = \frac{1}{2} BD$

$AC \perp BD$

To Prove: $4AB^2 = AC^2 + BD^2$

Proof: $\angle AOB = 90^\circ$

(Diagonals of rhombus bisect each other at right angle)

$$AB^2 = OA^2 + OB^2$$

or $AB^2 = \left(\frac{AC}{2}\right)^2 + \left(\frac{BD}{2}\right)^2 \quad [1]$

$$AB^2 = \frac{AC^2}{4} + \frac{BD^2}{4}$$

$$4AB^2 = AC^2 + BD^2 \quad [1]$$


Hence proved

OR

Let AB be the building and CB be the ladder.

Then,

$AB = 24 \text{ m}$, $CB = 25 \text{ m}$ and $\angle CAB = 90^\circ$.



By Pythagoras' theorem, we have:

$$CB^2 = AB^2 + AC^2$$

$$\Rightarrow AC^2 = (CB^2 - AB^2) = [(25)^2 - (24)^2] \text{ m}^2 \\ = (625 - 576) \text{ m}^2 = 49 \text{ m}^2$$

$$\Rightarrow AC = \sqrt{49} \text{ m} = 7 \text{ m.}$$


Hence, the distance of the foot of the ladder from the building.

37. Step of Construction:

(i) Construct a $\triangle ABC$ in which $BC = 6 \text{ cm}$, $AB = 5 \text{ cm}$ and $\angle ABC = 60^\circ$.

(ii) Draw a ray BX such that $\angle CBX$ acute angle.

(iii) Locate 4 points B_1, B_2, B_3 and B_4 such that $BB_1 = B_1B_2 = B_2B_3 = B_3B_4$.




- (iv) Join B_4C .
 (v) Through B_3 draw a line parallel to B_4C which meet BC at C' .
 (vi) Through C draw a line parallel to AC which meet AB at A' .
 (vii) $\triangle A'BC'$ is the required triangle.

[2 + 2]

38. Let AB be the building and CD be the tower such that $\angle BDE = 30^\circ$, $\angle BCA = 60^\circ$ and $AB = 60$ m.

Let $CA = DE = x$ metres.



From right $\triangle CAB$, we have:

$$\begin{aligned} \frac{CA}{AB} &= \cot 60^\circ = \frac{1}{\sqrt{3}} \\ \Rightarrow \frac{x}{60} &= \frac{1}{\sqrt{3}} \quad [1] \\ \Rightarrow x &= \left(60 \times \frac{1}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} \right) = 20\sqrt{3} \\ \Rightarrow CA &= DE = 20\sqrt{3} \text{ m} \quad \dots (i) [1] \end{aligned}$$

From right $\triangle BED$, we have:

$$\Rightarrow \frac{BE}{DE} = \tan 30^\circ = \frac{1}{\sqrt{3}} \Rightarrow \frac{BE}{20\sqrt{3}} = \frac{1}{\sqrt{3}} \quad [\text{using (i)}]$$

$$\Rightarrow BE = \left(20\sqrt{3} \times \frac{1}{\sqrt{3}} \right) = 20 \text{ m} \quad [1]$$

$$\therefore CD = AE = (AB - BE) \\ = (60 - 20) \text{ m} = 40 \text{ m.} \quad [1]$$

Hence, the height of the tower 40 m.

39. Given, Height of cylinder = 15 cm
 Its diameter = 12 cm
 radius = 6 cm
 radius of cone = 3 cm
 and height of cone = 9 cm [1]
 Let the number of toys recast be n .
 \therefore Volume of n conical toys = Volume of cylinder [1]

$$\Rightarrow n \times \frac{1}{3} \pi \times 3 \times 3 \times 9 = \pi \times 6 \times 6 \times 15$$

$$\Rightarrow n = \frac{6 \times 6 \times 15}{3 \times 9} \quad [1]$$

$$\Rightarrow n = 20 \quad [1]$$

Hence, the number of toys = 20

40.

| Students | c.f. |
|--------------|------|
| Less than 7 | 20 |
| Less than 9 | 38 |
| Less than 11 | 60 |
| Less than 13 | 85 |
| Less than 15 | 105 |
| Less than 17 | 120 |
| Less than 19 | 130 |

[1]

This curve is the required cumulative frequency curve or an ogive of the less than type.


Here, $N = 130$

$$\text{So, } \frac{N}{2} = \frac{130}{2} = 65 \quad [1]$$

Now, we locate the point on the ogive whose ordinate is 65. The x -co-ordinate corresponding to this ordinate is 11.4

Hence, the required median on the graph is 11.4. [1]

Units on: x -axis 1 cm = 2
 x -axis 1 cm = 10



[1]

OR

$$\text{Modal class} = 11 - 13$$

$$l = 11, f_1 = 95, f_0 = 41, f_2 = 36, h = 2$$

$$\text{Mode} = l + \frac{f_1 - f_0}{2f_1 - f_0 - f_2} \times 2$$

$$= 11 + \frac{95 - 41}{190 - 41 - 36} \times 2$$

$$= 11 + \frac{54}{113} \times 2$$

$$\text{Mode} = 11 + 0.95 = 11.95$$